

Beitr. Ent.	Keltern	ISSN 0005 - 805X
51 (2001) 1	S. 231 - 245	14.09.2001

Compilation of canthariphilous insects

CLAUDIA HEMP and KONRAD DETTNER

Summary

A list of insects attracted to cantharidin is given. Most canthariphilous insects are found within the hetero-meran beetle family Anthicidae with 190 species in the three subfamilies Anthicinae (184), Lemodiinae (1) and Tomoderinae (5). Further cantharidin baited species are known from the beetle families Endomychidae (5), Cleridae (4), Chrysomelidae (3), and Staphylinidae (1). In the beetle family Pyrochroidae 23 species from the subfamily Pedilinae and 6 species of the Pyrochroinae are canthariphilous. The insect order Diptera is represented by the families Anthomyiidae (3), Cecidomyiidae (2), Ceratopogonidae (22) Chloropidae (1), Platystomatidae (1) and Sciaridae (5). Many species from the heteropteran family Miridae (29) are known to be attracted by cantharidin, also one species from the families Lygaeidae and Tingidae respectively. Parasites of the hymenopteran family Braconidae (6) show a positive reaction to cantharidin as well as species from the subfamily Diapriinae of the family Diapriidae. Chemical analyses of a fulgorid and a cicadid species revealed that also species of Homoptera may contain cantharidin.

Keywords

cantharidin; canthariphilous insects; Coleoptera; Diptera; Heteroptera; Homoptera

Zusammenfassung

Die durch Cantharidin angezogenen Insekten werden aufgelistet. Die meisten Cantharidin-abhängigen Insekten sind in der Käferfamilie Anthicidae mit 190 Arten in den drei Unterfamilien Anthicinae (184), Lemodiinae (1) und Tomoderinae (5) gefunden worden. Weitere Cantharidin-abhängige Arten sind bekannt aus den Käferfamilien Endomychidae (5), Cleridae (4), Chrysomelidae (3) und Staphylinidae (1). In der Familie Pyrochroidae sind 23 Arten aus der Unterfamilie Pedilinae und 6 Arten aus den Pyrochroinae Cantharidin-beeinflusst. Bei den Dipteren sind dies in den Familien Anthomyiidae (3), Cecidomyiidae (2), Ceratopogonidae (22), Chloropidae (1), Platystomatidae (1) und Sciaridae (5). Manche Arten der Heteropterengruppe Miridae (29) sind bekannt wegen des Einflusses von Cantharidin ebenso jeweils eine Art aus den Familien Lygaeidae und Tingidae. Parasiten der Hymenopterengruppe Braconidae (6) zeigen eine positive Reaktion auf Cantharidin, ebenso wie Arten der Unterfamilie Diapriinae der Familie Diapriidae. Chemische Analysen von Arten der Fulgoridae und Cicadidae zeigen, dass auch Arten der Homoptera Cantharidin-abhängig scheinen.

Introduction

Cantharidin, a formal monoterpene anhydride, is known to humans for its toxic properties for over 2000 years. Applied on the human skin it induces serum filled blisters, which gave blister beetles (Coleoptera: Meloidae) their common name, beside Oedemeridae the only known natural sources of this chemical. Used internally cantharidin acts as diuretic and abortifacient. It may induce priapism, which made cantharidin one of the most well known aphrodisiacs.

In meloid and oedemerid beetles cantharidin is used as an effective haemolymph poison (CAVILL & CLARK 1971, CAPINERA et al. 1985, CARREL et al. 1986, BLODGETT et al. 1991, NICHOLLS et al. 1990, HOLZ et al. 1994), which protects the adults as well as larval stages and eggs.

Another interesting phenomenon in connection with cantharidin is its attractancy on other insects, which perceive this insecticide partly over great distances (GÖRNITZ 1937, WIRTH 1980, YOUNG 1984 a, b). Many of these so-called canthariphilous insects ingest cantharidin without any obvious damage.

This paper gives a compilation of insects attracted to cantharidin on species level. Canthariphilous species are known from following orders and families: Coleoptera (Anthicidae, Cleridae, Chrysomelidae, Endomychidae, Pyrochroidae, Staphylinidae), Diptera (Anthomyiidae, Cecidomyiidae, Ceratopogonidae, Chloropidae, Platystomatidae, Sciaridae), Heteroptera (Miridae, Lygaeidae, Tingidae), and Hymenoptera (Braconidae, Diapriidae). Analysis of Homoptera species proved, that cantharidin containing species can also be found in these order of insects.

Possible function of cantharidin in the life cycles of canthariphilous insects

Coleoptera

Most canthariphilous species are found in the beetle family Anthicidae with 190 recorded species. Mostly *Notoxus* and *Aulacoderus* species are attracted. Conspicuous in many anthicid genera associated with cantharidin are notches at the tips of the elytra in males which function as test organs. Females bite into these structures and choose their sexual partner by the amount of cantharidin a male has taken up (SCHÜTZ & DETTNER 1992). Ingested cantharidin is stored in the reproductive organs of the males and transmitted to females during copulation. The nuptial gift is secreted into eggs and thus provides protection for the offspring.

From the family Cleridae *Pallenothriocera rufimembris* is attracted to the terpenoid cantharidin (HEMP et al. 1999b). Also single specimens of other clerid species were collected in cantharidin traps put out in southern Europe. Further clerids were suggested to be canthariphilous (CHANDLER 1976, BOLOGNA & HAVELK 1984) or reported to contain cantharidin (JUANJIE et al. 1995, DETTNER, in prep.). Gland-like structures on the elytra of male *Pallenothriocera rufimembris* suggests an analogous cycle as found in anthicids. Specimens of a up to now unidentified clerid species attracted to cantharidin baits near Nairobi, Kenya were observed to remain several hours in the open traps, attacking there other arriving canthariphilous species (mostly *Notoxus* sp.), killing and feeding on them. Remaining elytra of the preys showed that one clerid often fed more than 10 *Notoxus* during the day hours.

Canthariphilous pyrochroid beetles show a similar cycle for cantharidin as discussed above for anthicids (HOLZ et al. 1994, EISNER et al. 1996b). Test organs for cantharidin titres are present on the head of the male in form of grooves (EISNER et al. 1996a). As in anthicid beetles the females bite into these structures to inform themselves about the amount of cantharidin the would-be partner has taken up.

For all other beetles families (Chrysomelidae, Endomychidae, Staphylinidae) with canthariphilous species only the phenomenon that members are attracted to the natural product cantharidin is known. No investigations have been undertaken up to now to illuminate life cycles and function of cantharidin for these insects.

Diptera and Heteroptera

Ceratopogonids are an ever present group as soon as cantharidin is put out. Studies on the function of cantharidin in the biology of canthariphilous ceratopogonids have been made by FRENZEL & DETTNER (1994). Both sexes of the investigated canthariphilous gnats are attracted to cantharidin, ingesting the terpenoid readily. Maximal concentrations in tissues of ceratopogonids analysed are similar to those detected in the cantharidin producing species of the heteromeren beetle families Meloidae and Oedemeridae. High concentrations of cantharidin in the haemolymph of European ceratopogonids proved to deter Empididae, which prey on these gnats.

As for most beetle families only the phenomenon that dipterans and heteropterans from families like Anthomyiidae, Cecidomyiidae, Chloropidae, Platystomatidae, Sciaridae (Diptera) and Miridae, Lygaeidae and Tingidae (Heteroptera) are attracted, is noted.

Hymenoptera

Braconids are parasitoids of other insects. Females are capable of sensing their host for egg-deposition via kairomones. This may also be the case in canthariphilous species that are attracted by the cantharidin smell of their hosts as suggested for *Perilissus plumicornis* which is a parasite of the anthicid *Notoxus monoceros* (GÖRNITZ 1937).

List of insects attracted to cantharidin

Coleoptera

Anthicidae
Anthicinae

<i>Acanthinus scitulus</i> (LECONTE)	CHANDLER 1976, YOUNG 1984a, b
<i>Anthicus lutulentus</i> CASEY	CHANDLER 1976, YOUNG 1984a
<i>Anthicus nanus</i> LECONTE	CHANDLER 1976, YOUNG 1984a
<i>Anthicus punctulatus</i> LECONTE	YOUNG 1984b
<i>Anthicus sonoranus</i> WERNER	YOUNG 1984b
<i>Aulacoderus albatarsis</i> (LAFERTÉ) ¹	VAN HILLE 1954
<i>Aulacoderus apterus</i> (VAN HILLE)	VAN HILLE 1984a
<i>Aulacoderus asymmetricus</i> (VAN HILLE)	VAN HILLE 1984a
<i>Aulacoderus bicoloritarsis</i> (PIC) ²	VAN HILLE 1954

- Aulacoderus bilineatus* (VAN HILLE)
Aulacoderus bonadonai (VAN HILLE)
Aulacoderus bradfordi (VAN HILLE)
Aulacoderus brevicornis (VAN HILLE)
Aulacoderus bryanti (PIC)
Aulacoderus canariensis (WOLLASTON)
Aulacoderus canthariphilus (VAN HILLE)
Aulacoderus cecileae (VAN HILLE)
Aulacoderus chappuisi (PIC)
Aulacoderus citernii (PIC)
Aulacoderus colletti (VAN HILLE)
Aulacoderus dorsalis (LAFERTÉ)
Aulacoderus flavitarsis (FAHRAEUS)⁴
Aulacoderus flavopictus (LAFERTÉ)
Aulacoderus forchhammeri (VAN HILLE)
Aulacoderus formicomisternus (VAN HILLE)
Aulacoderus forsythi (VAN HILLE)
Aulacoderus fragilis (FAHRAEUS)
Aulacoderus govenderi (VAN HILLE)
Aulacoderus halleyi (VAN HILLE)
Aulacoderus inopinans (KREKICH)
Aulacoderus kochi (VAN HILLE)
Aulacoderus maccleryi (VAN HILLE)
Aulacoderus martini (PIC)
Aulacoderus mediofasciatus (PIC)
Aulacoderus milleri (VAN HILLE)
Aulacoderus mogotoensis (VAN HILLE)
Aulacoderus multidenticulatus (VAN HILLE)
Aulacoderus munroi (VAN HILLE)
Aulacoderus mutatus (GEMMINGER)
Aulacoderus orangensis (VAN HILLE)
Aulacoderus pedester (VAN HILLE)
Aulacoderus perlucidus (VAN HILLE)
Aulacoderus poweri (VAN HILLE)
Aulacoderus ranchbodi (VAN HILLE)
Aulacoderus recognitus (PIC)
Aulacoderus reverendus (PIC)
Aulacoderus rotundipennis (PIC)
Aulacoderus scydmaenoides (WOLLASTON)
Aulacoderus serowensis (VAN HILLE)
Aulacoderus sibayensis (VAN HILLE)
Aulacoderus simoni (PIC)⁷
Aulacoderus smithersi (VAN HILLE)
Aulacoderus techowi (PIC)
- Cordicomus instabilis* (SCHMIDT) pers. comm.⁹
- Cyclodinus basilewskyi* BUCK pers. comm.¹⁰
- VAN HILLE 1984a
FORCHHAMMER 1985
VAN HILLE 1984a
VAN HILLE 1984b
pers. comm.³
SCHÜTZ & DETTNER 1992
VAN HILLE 1984a
VAN HILLE 1985
HEMP et al. 1999a
VAN HILLE 1984a
pers. observ., 49
VAN HILLE 1984a
VAN HILLE 1984a
VAN HILLE 1984a
VAN HILLE 1984a
VAN HILLE 1954, 1984a
VAN HILLE 1985
VAN HILLE 1984a
VAN HILLE 1984a
VAN HILLE 1984a, pers. observ.⁵
VAN HILLE 1984b
VAN HILLE 1988
HEMP et al. 1999a
VAN HILLE 1985
VAN HILLE 1984a
HEMP 1994⁶
VAN HILLE 1985
VAN HILLE 1971, 1984a
VAN HILLE 1954, 1984a
VAN HILLE 1984a
VAN HILLE 1985, pers. observ.⁸

<i>Endomia tenuicollis</i> (ROSSI)	BOLOGNA & HAVELKA 1984
<i>Formicilla munda</i> (LECONTE)	YOUNG 1984b
<i>Formicomus caeruleus</i> (THUNBERG)	VAN HILLE 1954, 1985
<i>Formicomus canaliculatus</i> LAFERTÉ	VAN HILLE 1971, HEMP et al. 1999a
<i>Formicomus chappuisi</i> PIC	pers. observ. ¹¹
<i>Formicomus consul</i> LAFERTÉ	YOUNG 1984b
<i>Formicomus gestroi</i> PIC	SCHÜTZ & DETTNER 1992, HEMP et al. 1997, HEMP & DETTNER 1997, HEMP et al. 1999a
<i>Formicomus lacustris</i> KREK.	VAN HILLE 1971, HEMP et al. 1999a
<i>Formicomus linnavouri</i> VAN HILLE	pers. observ. ⁵⁰
<i>Formicomus millerianus</i> PIC	HEMP et al. 1999a
<i>Formicomus opaculus</i> KOLBE	HEMP et al. 1999a
<i>Formicomus pedestris</i> (ROSSI)	GÖRNITZ 1937, BOLOGNA & HAVELKA 1984, HEMP & DETTNER 1997
<i>Formicomus rubricollis</i> LAFERTÉ	SCHÜTZ & DETTNER 1992, HEMP et al. 1997, HEMP & DETTNER 1997, HEMP et al. 1999a
<i>Formicomus</i> sp. (<i>lewisi</i> MARSEUL?)	YOUNG 1984b
<i>Formicomus spatulatus</i> VAN HILLE	HEMP et al. 1999a
<i>Formicomus tropicalis</i> KREK.	VAN HILLE 1971
<i>Hirticomus biplagiatus</i> (LAFERTÉ)	pers. observ. ⁵¹
<i>Hirticomus hispidus</i> (ROSSI)	pers. observ. ¹²
<i>Hirticomus quadriguttatus</i> (ROSSI)	BOLOGNA & HAVELKA 1984, pers. observ. ¹³
<i>Mecynotarsus balsensis</i> WERNER	CHANDLER 1976, 1977b
<i>Mecynotarsus casperi</i> PIC	pers. comm. ¹⁴
<i>Mecynotarsus falcatus</i> CHANDLER	YOUNG 1984b
<i>Mecynotarsus lacustris</i> VAN HILLE	VAN HILLE 1971
<i>Mecynotarsus nevermanni</i> WERNER	HEMP et al. 1999a
<i>Mecynotarsus nigronotatus</i> PIC	HEMP et al. 1999a
<i>Mecynotarsus obliquemaculatus</i> MARSEUL	YOUNG 1984b
<i>Mecynotarsus oblitteratus</i> PIC	HEMP et al. 1999a
<i>Mecynotarsus serricornis</i> (PANZER)	BOLOGNA & HAVELKA 1984
<i>Mecynotarsus vagipictus</i> FAIRMAIRE	YOUNG 1984b
<i>Microhoria aspelia</i> (TRUQUI)	pers. comm. ¹⁵
<i>Microhoria aubei</i> (LAFERTÉ) ¹⁶	GÖRNITZ 1937, ABDULLAH 1964, YOUNG 1984a
<i>Microhoria biauriculatus</i> (PIC) ¹⁷	GÖRNITZ 1937, ABDULLAH 1964, YOUNG 1984a
<i>Microhoria calliger</i> (MARSEUL)	pers. comm. ¹⁸
<i>Microhoria chobauti</i> (PIC) ¹⁹	GÖRNITZ 1937, ABDULLAH 1964, YOUNG 1984a
<i>Microhoria cinctuta</i> (MARSEUL) ²⁰	GÖRNITZ 1937, ABDULLAH 1964, BOLOGNA & HAVELKA 1984, YOUNG 1984a
<i>Microhoria fairmairei</i> (BRISOUT) ²¹	GÖRNITZ 1937, ABDULLAH 1964, YOUNG 1984a
<i>Microhoria fasciata</i> (CHEVROLAT)	BOLOGNA & HAVELKA 1984
<i>Microhoria fasciata</i> (CHEVROLAT) <i>codinai</i> PIC	BOLOGNA & HAVELKA 1984

<i>Microboria ghilianii</i> (LAFERTÉ)	BOLOGNA & HAVELKA 1984
<i>Microboria insignis</i> (LUCAS) ²²	GÖRNITZ 1937, YOUNG 1984a
<i>Microboria insignis</i> (LUCAS) var. <i>panousei</i> (PIC) ²³	YOUNG 1984a
<i>Microboria obscuripes</i> (PIC) ²⁴	ABDULLAH, 1964, YOUNG 1984a
<i>Microboria plagifer</i> (KREKICH)	pers. observ. ²⁵
<i>Microboria syrensis</i> (PIC)	pers. comm. ²⁶
<i>Microboria terminata</i> (SCHMIDT)	SCHÜTZ & DETTNER 1992, HEMP & DETTNER 1997
<i>Microboria tortiscelis</i> (MARSEUL) ²⁷	YOUNG 1984a
<i>Microboria venusta</i> (VILLA)	BOLOGNA & HAVELKA 1984
<i>Notoxus allansonii</i> VAN HILLE	VAN HILLE 1971
<i>Notoxus alluaudi</i> PIC	HEMP et al. 1999a
<i>Notoxus amaculatus</i> VAN HILLE	VAN HILLE 1971
<i>Notoxus anchora</i> HENTZ	CHANDLER 1977a, 1982
<i>Notoxus bifasciatus</i> (LECONTE)	YOUNG 1984b
<i>Notoxus brutoni</i> VAN HILLE	VAN HILLE 1985
<i>Notoxus buraensis</i> UHMANN	HEMP et al. 1999a
<i>Notoxus calcaratus</i> HORN	CHANDLER 1976, 1982, YOUNG 1984a
<i>Notoxus candatus</i> FALL	CHANDLER 1982
<i>Notoxus cavicornis</i> LECONTE	CHANDLER 1982
<i>Notoxus celatus</i> CHANDLER	CHANDLER 1977a
<i>Notoxus conformis</i> LECONTE	CHANDLER 1977a, 1982
<i>Notoxus constrictus</i> CASEY	CHANDLER 1977a, YOUNG 1984a
<i>Notoxus cornutus</i> THUNBERG	ABDULLAH 1964
<i>Notoxus cucullatus</i> LAFERTÉ	VAN HILLE 1971
<i>Notoxus cumanensis</i> LAFERTÉ	CHANDLER 1977a
<i>Notoxus daressalaamensis</i> UHMANN	HEMP et al. 1999a
<i>Notoxus decorus</i> VAN HILLE	SCHÜTZ & DETTNER 1992, HEMP et al. 1999a
<i>Notoxus denudatus</i> HORN	CHANDLER 1982
<i>Notoxus desertus</i> CASEY	CHANDLER 1977a, 1982, YOUNG 1984a
<i>Notoxus filicornis</i> CASEY	YOUNG 1984b
<i>Notoxus fraternus</i> CHAMPION	CHANDLER 1977a
<i>Notoxus guttulatus</i> BUCK	ABDULLAH 1964, pers. comm. ²⁸
<i>Notoxus hageni</i> CHANDLER	CHANDLER 1982
<i>Notoxus hiltoni</i> VAN HILLE	VAN HILLE 1971
<i>Notoxus hirsutus</i> CHAMPION	CHANDLER 1977a, 1982
<i>Notoxus hirsutus</i> PIC	pers. observ. ⁵²
<i>Notoxus hirtus</i> LAFERTÉ	BOLOGNA & HAVELKA 1984
<i>Notoxus holmi</i> UHMANN	VAN HILLE 1985
<i>Notoxus intermedius</i> FALL	CHANDLER 1982
<i>Notoxus lateralis</i> CHANDLER	CHANDLER 1977a
<i>Notoxus lesnei</i> PIC	pers. comm. ²⁹
<i>Notoxus lunulifer</i> PIC	VAN HILLE 1971, HEMP et al. 1999a
<i>Notoxus marginatus</i> LECONTE	CHANDLER 1976, 1977a, 1982
<i>Notoxus mauritanicus</i> LAFERTÉ	ABDULLAH 1964, YOUNG 1984a
<i>Notoxus mexicanus</i> CHAMPION	CHANDLER 1976, 1977a
<i>Notoxus mkuziensis</i> VAN HILLE	VAN HILLE 1971

<i>Notoxus monoceros</i> (LINNÉ)	GÖRNITZ 1937, ABDULLAH 1964, YOUNG 1984a, BOLOGNA & HAVELKA 1984
<i>Notoxus monodon</i> (FABRICIUS)	CHANDLER 1976, 1982
<i>Notoxus montanus</i> CASEY	CHANDLER 1982
<i>Notoxus murinipennis</i> (LECONTE)	CHANDLER 1976, 1977a, 1982
<i>Notoxus namibianus</i> UHMANN	pers. observ. ³⁰
<i>Notoxus nevadensis</i> CASEY	CHANDLER 1982
<i>Notoxus nuperus</i> HORN <i>hastrus</i> CHANDLER	CHANDLER 1982
<i>Notoxus nuperus</i> HORN	CHANDLER 1976
<i>Notoxus opacus</i> CHAMPION	CHANDLER 1977a
<i>Notoxus ornatus</i> VAN HILLE	VAN HILLE 1971
<i>Notoxus photus</i> CHANDLER	YOUNG 1984b
<i>Notoxus pictus</i> CASEY	CHANDLER 1982
<i>Notoxus planicornis</i> LAFERTÉ	pers. comm. ³¹
<i>Notoxus pretiosus</i> VAN HILLE	HEMP et al. 1999a
<i>Notoxus pygidialis</i> CHANDLER	CHANDLER 1977a
<i>Notoxus reavelli</i> VAN HILLE	VAN HILLE 1984b
<i>Notoxus recticornis</i> VAN HILLE	pers. comm. ³²
<i>Notoxus robustus</i> CASEY	CHANDLER 1982, YOUNG 1984a
<i>Notoxus roehri</i> UHMANN	VAN HILLE 1985, pers. comm. ³³
<i>Notoxus rothschildi</i> PIC	HEMP et al. 1999a
<i>Notoxus rufomaculatus</i> PIC	HEMP et al. 1999a
<i>Notoxus seminole</i> CHANDLER	YOUNG 1984a, b
<i>Notoxus serratus</i> (LECONTE)	CHANDLER 1977a, 1982
<i>Notoxus sparsus</i> LECONTE	CHANDLER 1977a, 1982, YOUNG 1984a
<i>Notoxus spatulifer</i> CASEY	CHANDLER 1982
<i>Notoxus talpa</i> LAFERTÉ	CHANDLER 1976, 1977a
<i>Notoxus tansanianus</i> UHMANN	HEMP et al. 1999a
<i>Notoxus toltecorum</i> CHANDLER	CHANDLER 1977a
<i>Notoxus vanhillei</i> UHMANN	HEMP et al. 1999a
<i>Notoxus whartoni</i> CHANDLER	CHANDLER 1982
<i>Notoxus walteri</i> VAN HILLE	pers. observ. ⁵³
<i>Notoxus youngi</i> CHANDLER	CHANDLER 1982
<i>Notoxus zapotecorum</i> CHANDLER	CHANDLER 1977a
<i>Omonadus bottegoi</i> PIC	pers. comm. ³⁴
<i>Pseudoleptaleus unifasciatus</i> (DESBR.)	FORCHHAMMER 1985, HEMP et al. 1999a
<i>Pseudonotoxus testaceus</i> (LAFERTÉ)	VAN HILLE 1985
<i>Sapintus javanus</i> (MARSEUL)	YOUNG 1984b, pers. comm. ³⁵
<i>Sapintus malayensis</i> (PIC) ³⁶	GÖRNITZ 1937
<i>Sapintus plectilis</i> (PIC)	YOUNG 1984b
<i>Sapintus pollocki</i> UHMANN	pers. comm. ³⁷
<i>Sapintus tavetanus</i> (PIC)	HEMP et al. 1999a
<i>Tenuicomus babaulti</i> (PIC)	HEMP et al. 1999a

<i>Tenuicomus barnevillei</i> (PIC)	BOLOGNA & HAVELKA 1984
<i>Tenuicomus pumilus</i> (BAUDI) ³⁸	GÖRNITZ 1937, ABDULLAH 1964, YOUNG 1984a
<i>Vacusus infernus</i> (LAFERTÉ)	CHANDLER 1976
 Lemodiinae	
<i>Trichanana aptera</i> LEA	pers. observ. ³⁹
 Tomoderinae	
<i>Tomoderus</i> sp.	CHANDLER 1976, YOUNG 1984b
<i>Tomoderus alluaudi</i> PIC	HEMP et al. 1999a
<i>Tomoderus brevicornis</i> CHAMPION	pers. observ. ⁴⁰
<i>Tomoderus congoanus</i> PIC	HEMP et al. 1999a
<i>Tomoderus kolbei</i> PIC	HEMP et al. 1999a
 Endomychidae	
<i>Aphorista laeta</i> (LECONTE)	YOUNG 1984b
<i>Aphorista vittata</i> (FABRICIUS)	YOUNG 1984b
 <i>Danae testacea</i> (ZIEGLER)	YOUNG 1984b
 <i>Lycoperdina ferruginea</i> LECONTE	YOUNG 1984b
 <i>Xenomycetes morrisoni</i> HORN	YOUNG 1989, DETTNER 1997
 Cleridae	
<i>Cymatodera</i> sp.	CHANDLER 1976
 <i>Pallenothriocera rufimembris</i> PIC	HEMP et al. 1999a, HEMP et al. 1999b
 Gen. sp.	BOLOGNA & HAVELKA 1984
 Cleridae sp.	pers. observ. ⁴¹
 Chrysomelidae	
<i>Aristobrotica angulicollis</i> (ERICHSON)	MAFRA-NETO & JOLIVET 1994
 <i>Barombiella</i> (Bonesioides) <i>vicina</i> LABOISSIERE	HEMP et al. 1999a
<i>Barombiella</i> (Bonesioides) ssp.	HEMP et al. 1999a
 Pyrochroidae	
 Pedilinae	
<i>Anisotria shooki</i> YOUNG	YOUNG 1984a
 <i>Pedilus abnormis</i> (HORN)	YOUNG 1981
<i>Pedilus bardi</i> (HORN)	YOUNG 1981
<i>Pedilus canaliculatus</i> (LECONTE)	YOUNG 1981

<i>Pedilus cavatus</i> FALL	YOUNG 1981
<i>Pedilus collaris</i> (SAY)	BUTLER 1984
<i>Pedilus crotchi</i> (HORN)	YOUNG 1981
<i>Pedilus elegans</i> (HENTZ)	YOUNG 1981, YOUNG 1984a
<i>Pedilus impressus</i> (SAY)	ABDULLAH 1964, YOUNG 1981, YOUNG 1984a
<i>Pedilus inconspicuus</i> (HORN)	YOUNG 1981
<i>Pedilus joanae</i> YOUNG	YOUNG 1981
<i>Pedilus johnsonorum</i> YOUNG	YOUNG 1981
<i>Pedilus labiatus</i> (SAY)	ABDULLAH 1964, YOUNG 1981
<i>Pedilus lewisi</i> (HORN)	YOUNG 1981
<i>Pedilus longlobus</i> FALL	YOUNG 1981
<i>Pedilus lugubris</i> (SAY)	YOUNG 1981, YOUNG 1984a
<i>Pedilus monticolus</i> (HORN)	YOUNG 1981
<i>Pedilus oregonus</i> FALL	YOUNG 1981
<i>Pedilus picipennis</i> FALL	YOUNG 1981
<i>Pedilus punctulatus</i> LECONTE	YOUNG 1981
<i>Pedilus serratus</i> FALL	YOUNG 1981
<i>Pedilus</i> sp. nov.	YOUNG 1984a
<i>Pedilus terminalis</i> (SAY)	YOUNG 1981, YOUNG 1984a
Pyrochroinae	
<i>Neopyrochroa femoralis</i> (LECONTE)	YOUNG 1984b
<i>Neopyrochroa flabellata</i> (FABRICIUS)	YOUNG 1984a
<i>Pyrochroa coccinea</i> LINNÉ	HOLZ et al. 1994
<i>Pyrochroa serraticornis</i> (SCOPOLI)	BOLOGNA & HAVELKA 1984
<i>Schizotus cervicalis</i> NEWMANN	YOUNG 1984b
<i>Schizotus pectinicornis</i> LINNÉ	HOLZ et al. 1994
Staphylinidae	
<i>Eusphalerum minutum</i> (F.)	DETTNER 1997
Diptera	
Anthomyiidae	
<i>Anthomyia benguellae</i> MALLOCH	HEMP et al. 1999a
<i>Anthomyia pluvialis</i> (LINNÉ)	GÖRNITZ 1937, BOLOGNA & HAVELKA 1984, FRENZEL & DETTNER 1994
<i>Delia (Hylemyia) trispinosa</i> KARL	GÖRNITZ 1937, DETTNER 1997
Cecidomyiidae	
Gen. sp. Cecidomyiinae	DETTNER 1997
Gen. sp. Lestremiinae	DETTNER 1997
Ceratopogonidae	
<i>Atrichopogon brunnipes</i> MEIGEN	GÖRNITZ 1937 ⁴² , FRENZEL et al. 1992
<i>Atrichopogon downesi</i> WIRTH	WIRTH 1980

<i>Atrichopogon epicautae</i> WIRTH	WIRTH 1980
<i>Atrichopogon favri</i> WIRTH	WIRTH 1980
<i>Atrichopogon geminus</i> BOESEL	YOUNG 1984b
<i>Atrichopogon illiesi</i> HAVELKA	BOLOGNA & HAVELKA 1984
<i>Atrichopogon levius</i> (COQUILLET)	YOUNG 1984b
<i>Atrichopogon lindneri</i> WIRTH	WIRTH 1980
<i>Atrichopogon lucorum</i> (MEIGEN)	BOLOGNA & HAVELKA 1984, FRENZEL et al. 1992
<i>Atrichopogon maculosus</i> EWEN	YOUNG 1984b
<i>Atrichopogon meloesugans</i> KIEFFER	WIRTH 1980, BOLOGNA & HAVELKA 1984
<i>Atrichopogon mittmanni</i> HAVELKA	pers. observ. ⁴³
<i>Atrichopogon oedemerarum</i> STORA	FRENZEL et al. 1992, FRENZEL & DETTNER 1994
<i>Atrichopogon rostratus</i> (WINNERTZ)	WIRTH 1956
<i>Atrichopogon ruediger</i> HAVELKA	pers. observ. ⁴⁴
<i>Atrichopogon setospennis</i> KIEFFER	BOLOGNA & HAVELKA 1984
<i>Atrichopogon</i> sp.	BOLOGNA & HAVELKA 1984
<i>Atrichopogon</i> sp. near <i>websteri</i> (COQUILLET)	YOUNG 1984b
<i>Atrichopogon trifasciatus</i> KIEFFER	BOLOGNA & HAVELKA 1984, FRENZEL et al. 1992, FRENZEL & DETTNER 1994
<i>Atrichopogon websteri</i> (COQUILLET)	YOUNG 1984b
<i>Culicoides obsoletus</i> MEIGEN	BOLOGNA & HAVELKA 1984
<i>Forcipomyia crinita</i> SAUNDERS	BOLOGNA & HAVELKA 1984
Chloropidae	
<i>Goniopsita</i> spec.	DETTNER 1997
Platystomatidae	
<i>Peltacanthina</i> near <i>mythodes</i> HENDEL	HEMP et al. 1999a
Sciaridae	
ssp.	YOUNG 1984b
Gen. sp.	DETTNER 1997 ⁴⁵
<i>Bradysia optata</i>	RUDZINSKI, FRANK & DETTNER, 2001
<i>Heterosciara</i> sp.	pers. observ. ⁴⁶
<i>Schwenkfeldina</i> sp.	pers. observ. ⁴⁷
<i>Scythropochroa</i> sp.	pers. observ. ⁴⁸

Heteroptera

Lygaeidae

Dieuches sp.HEMP & DETTNER, in press, HEMP et al.
1999a

Miridae

Caulotops ssp.

YOUNG 1984b

Cryptocapsus paraensis (CARVALHO)*Cryptocapsus* sp.

MAFRA-NETO & JOLIVET 1994

DETTNER 1997

Cryptocapsus caligineus (STAL)

HEMP & DETTNER, 2000

Erychilella sp. near *pallida* REUTER

YOUNG 1984b

Hadroneura bispinosa KNIGHT

PINTO 1978

Hadroneura brevifata KNIGHT

YOUNG 1984b

Hadroneura militaris UHLER

FOX 1943

Hadroneura princeps UHLER

YOUNG 1984b

Hadroneura sp.

PINTO 1978

Hadroneura ulbieri VANDUZEE

PINTO 1978, YOUNG 1984b

Hadroneura uniformis KNIGHT

YOUNG 1984b

Halticotoma nicholi KNIGHT

PINTO 1978

Halticotoma sp.

YOUNG 1984b

Halticotoma valida TOWNSEND

YOUNG 1984b

Neoleucon sulinus (CARVALHO)

MAFRA-NETO & JOLIVET 1994

Pachymerocista pilosus (CARVALHO)

HEMP & DETTNER, 2000

Pycnoderes quadrimaculatus (GUERIN-MELNEVILLE)

YOUNG 1984b, HEMP & DETTNER, 2000

Pycnoderes sp.

YOUNG 1984b

Pycnoderes vanduzeei REUTER

HEMP & DETTNER, 2000

Sixeonotus brevirostris KNIGHT

YOUNG 1984b

Sixeonotus insignis REUTER

YOUNG 1984b

Sixeonotus sp.

PINTO 1978, YOUNG 1984b, HEMP &

Dettner, 2000

DETTNER, 2000

Sixeonotus sp. near *brevirostris* KNIGHT

YOUNG 1984b

Sixeonotus tenebrosus (DISTANT)

YOUNG 1984b

Syninas fulvicollis (FABRICIUS)

MAFRA-NETO & JOLIVET 1994

Syninas linearis DISTANT

YOUNG 1984b

Syninas signaticollis (REUTER)

MAFRA-NETO & JOLIVET 1994

Thentecoris sp.

DETTNER 1997

Tingidae
sp.

MAFRÀ-NETO & JOLIVET 1994, DETTNER
1997

Hymenoptera

Braconidae

Blacus ruficornis (NEES)

YOUNG 1984b

Perilissus plumicornis RUTHE

GÖRNITZ 1937

Streblocera near *pulvillicornis* WALLEY &

MACKAY

YOUNG 1984b

Syrrhizus agilis (CRESSON)

YOUNG 1984b

Microtonus sp.

DETTNER 1997

Melittobia sp.

BOLOGNA 1991

Diapriidae

Gen. sp. (Diapriinae)

DETTNER 1997

¹ *Anthicus albitarvis* in VAN HILLE 1954

² *Anthicus bicoloritarsis* in VAN HILLE 1954

³ collected by Dr. POLLOCK at cantharidin in Australia (pers. comm UHMANN)

⁴ *Anthicus flavopictus* in VAN HILLE 1954 and 1984a

⁵ Expedition of the Museum Berlin to southern Africa (pers. comm UHMANN)

⁶ attracted to cantharidin on the canary island Tenerife

⁷ *Athicus simoni* in VAN HILLE 1954 and 1984a

⁸ Expedition of the Museum Berlin to southern Africa (pers. comm UHMANN)

⁹ collected by Dr. SCHAWALLER in Nepal, feeding on meloids

¹⁰ collected by Dr. T. WAGNER at cantharidin in Uganda, East Africa

¹¹ attracted to cantharidin bait in March 1999 at the Tanzanian coast near Pangani

¹² attracted to cantharidin bait on the mediterranean island Mallorca 7.90

¹³ attracted to cantharidin at the mediterranean island Mallorca 7.90

¹⁴ Expedition of the Museum Berlin to southern Africa (pers. comm UHMANN)

¹⁵ collected at cantharidin by C. KOCH in Cyprus

¹⁶ *Anthicus aubei* in GÖRNITZ 1937, ABDULLAH 1964 and YOUNG 1984a

¹⁷ *Anthicus biauriculatus* in GÖRNITZ 1937, ABDULLAH 1964 and YOUNG 1984a

¹⁸ collected at cantharidin bait by Dr. W. ARENS in Greece, Olympia

¹⁹ *Anthicus chobauti* in GÖRNITZ 1937, ABDULLAH 1964 and YOUNG 1984a

²⁰ *Anthicus cinctutus* in GÖRNITZ 1937, ABDULLAH 1964, BOLOGNA & HAVELK 1984 and YOUNG 1984a

²¹ *Anthicus fairmairei* in GÖRNITZ 1937, ABDULLAH 1964 and YOUNG 1984a

²² *Anthicus insignis* in GÖRNITZ 1937 and YOUNG 1984a

²³ *Anthicus panousei* in YOUNG 1984a

²⁴ *Anthicus obscuripes* in ABDULLAH 1964 and YOUNG 1984a

²⁵ at cantharidin on the mediterranean island Mallorca, July 1990

²⁶ collected at cantharidin by C. KOCH in Cyprus

²⁷ *Anthicus tortisealis* in YOUNG 1984a

²⁸ Expedition of the Museum Berlin to southern Africa (pers. comm UHMANN)

²⁹ Expedition of the Museum Berlin to southern Africa (pers. comm UHMANN)

- ³⁰ Expedition of the Museum Berlin to southern Africa (pers. comm UHMANN)
- ³¹ personal communication with Prof. CHANDLER, USA
- ³² around 150 specimens at cantharidin in December 1998 and April 1999 in the suburb Karen of Nairobi, Kenya
- ³³ Expedition of the Museum Berlin to southern Africa (pers. comm UHMANN)
- ³⁴ caught by Dr. T. WAGNER at cantharidin in Uganda, East Africa
- ³⁵ collected at cantharidin by Prof. CHANDLER on the Phillipines, Cebu, Inayagan and Naga
- ³⁶ *Anthicus malayensis* in GÖRNITZ 1937
- ³⁷ collected at cantharidin by Dr. POLLOCK, Australia (pers. comm UHMANN)
- ³⁸ *Anthicus pumilus* in GÖRNITZ 1937, ABDULLAH 1964 and YOUNG 1984a
- ³⁹ specimen labeled „at cantharidin“, coll. POLLOCK, Australia
- ⁴⁰ baited with cantharidin, southcoast, Dominican Republic, 1.95
- ⁴¹ in December 1998 about 50 specimens of a up to now not identified clerid species were attracted to cantharidin baits from a piece of indigenous *Olea*-forest at Karen near Nairobi, Kenya
- ⁴² *Kempia brunneipes* in GÖRNITZ 1937
- ⁴³ new species from Tanzania, Kilimanjaro, mentioned in FRENZEL et al. (1998), species description by HAVELKA, in prep.; listed in HEMP et al. 1999a under *Atrichopogon* spec. 1
- ⁴⁴ new species from Tanzania, Kilimanjaro, mentioned in FRENZEL et al. (1998), species description by HAVELKA, in prep.; listed in HEMP et al. 1999a, under *Atrichopogon* spec. 2
- ⁴⁵ 3 males were collected at Mistelbach/Bayreuth (August 90) at cantharidin
- ⁴⁶ 21 males and 2 females were collected at Bayreuth (June 1991) at cantharidin. According to MENZEL & MOHRIG (1997) *Heterosciara* is a synonym of *Scatopsiara*
- ⁴⁷ 1 male was collected at Bayreuth (June 1991) at cantharidin
- ⁴⁸ 4 males and 1 female were collected at Mistelbach/Bayreuth (August 1990) at cantharidin
- ⁴⁹ attracted to cantharidin in savannah grassland at 1400 m near Mt. Kitumbeine, Tanzania (11/99)
- ⁵⁰ attracted to cantharidin in coastal grassland near Pagani, Tanzania (1/00)
- ⁵¹ attracted to cantharidin in coastal grassland near Pagani, Tanzania (3/00)
- ⁵² attracted to cantharidin in savannah grassland near Mt. Kitumbeine, Tanzania (11/99)
- ⁵³ attracted to cantharidin in savannah grassland near Mt. Kitumbeine, Tanzania (11/99)

Homoptera

First results in analysing cantharidin contents in fulgorids suggest, that there are also members in the order Homoptera associated with cantharidin. The lantern fly *Lycorma delicatula* proved to contain cantharidin, which probably was taken up by host plants (FENG et al. 1988, DETTNER 1997). There was also suggested the Cicadidae species *Huechys sanguinea* (DE GEER) would contain cantharidin (JUANJIE et al. 1995).

Acknowledgements

We wish to thank all those persons who put out cantharidin for us to attract canthariphilous insects: Dr. W. ARENS, Bayreuth, C. KOCH, Bayreuth, Dr. T. WAGNER, Museum König Bonn, Dr. SCHAWALLER and Prof. CHANDLER who mentioned several canthariphilous species in our communication. Many thanks also to Mr. G. UHMANN, who kindly informed us about new „cases“ of canthariphily and for revising the manuscript.

References

- ABDULLAH, M. 1964: *Protomeloe argentinensis*, a new genus and species of Meloidae, with remarks on the significance of Cantharidin and the phylogeny of the families Pyrochroidae, Anthicidae, Meloidae and Cephaloidea. - Ann. Mag. Nat. Hist. 7: 247-254.
- BLODGETT, S. L.; CARREL, J. E. & HIGGINS, R. A. 1991: Cantharidin Content of Blister Beetles (Coleoptera: Meloidae) Collected from Kansas Alfalfa and Implications for Inducing Cantharidiasis. - Environmental Entomology 20 (3): 776-780.
- BOLOGNA, M. A. & HAVELKA, P. 1984: Nuove Segnalazioni Di Attrazione Della Cantharidina Dei Meloidae Su Coleotteri E Ditteri. - Boll. Assoc. Rom. Entomol. 39: 77-82.

- BOLOGNA, M. A. 1991: Fauna d'Italia. Coleoptera: Meloidae.
- CAPINERA, J. L.; GARDNER, D. R. & STERMITZ, F. R. 1985: Cantharidin levels in blister beetles (Coleoptera: Meloidae) associated with alfalfa in Colorado. - *J. Econ. Entomol.* **78**: 1052-1055.
- CARREL, J. E.; DOOM, J. P. & MCCORMICK, J. P. 1986: Identification of cantharidin in false blister beetles (Coleoptera, Oedemeridae) from Florida. - *J. Chem. Ecol.* **12**: 741-748.
- CAVILL, G. W. K. & CLARK, D. V. 1971: Ant secretions and cantharidin. In: M. JACOBSON, D. G. CROSBY (eds.). *Naturally Occurring Insecticides*. Marcel Dekker, New York: 271-304.
- CHANDLER, D. S. 1976: Use of Cantharidin and meloid beetles to attract Anthicidae (Coleoptera). - *Pan Pacific Entomol.* **52**: 179-80.
- CHANDLER, D. S. 1977a: A revision of the central and south american Notoxus and description of a new genus, Plesionotoxus. *Contr. - Am. Entomol. Inst.* **15**: 1-83.
- CHANDLER, D. S. 1977b: New Mecynotarsus with a key to the new world species (Coleoptera: Anthicidae). - *Col. Bull.* **31**: 363-370.
- DETTNER, K. 1997: Inter- and Intraspecific transfer of Toxic Insect Compound Cantharidin. In: DETTNER et al. (eds.). *Vertical Food Web Interactions*: 115-145.
- EISNER, T.; SMEDLEY, S. R.; YOUNG, D. K.; EISNER, M.; ROACH, B. & MEINWALD, J. 1996a: Chemical basis of courtship in a beetle (*Neopyrochroa flabellata*): Cantharidin as precopulatory „exciting“ agent. - *Proceedings of the National Academy of Sciences* **93** (13): 6494-6498.
- EISNER, T.; SMEDLEY, S. R.; YOUNG, D. K.; EISNER, M.; ROACH, B. & MEINWALD, J. 1996b: Chemical basis of courtship in a beetle (*Neopyrochroa flabellata*): Cantharidin as „nuptial gift“. - *Proceedings of the National Academy of Sciences* **93** (13): 6499-6498.
- FENG, Y.; JIANQI, M.; ZHONGREN L. & TIANPENG, G. 1988: A Preliminary Investigation on the Cantharidin Resources of Shaanxi Province. - *Acta Univ. Sept. Occident. Agric.* **16**: 23-28.
- FORCHHAMMER, P. 1985: Seasonal and daily variations in activity of ant-like flower-beetles (Anthicidae) collected in Serowe, Botswana 1982-83. - *Botswana Notes Rec.* **17**: 163-174.
- FOX, N. B. 1943: Some insects infesting the „selenium indicator“ vetches in Saskatchewan. - *Can. Entomol.* **75**: 206-207.
- FRANK, J. & DETTNER, K. 2001: Attraction of fungus gnat *Bradyia optata* to cantharidin. - *Entomol. Exp. Appl.*, in press
- FRENZEL, M.; DETTNER, K.; WIRTH, D.; WAIBEL, J. & BOLAND, W. 1992: Cantharidin analogues and their attractancy for ceratopogonid flies (Diptera: Ceratopogonidae). - *Experientia* **48**: 106-111.
- FRENZEL, M. & DETTNER, K. 1994: Quantitation of cantharidin in canthariphilous ceratopogonids (Diptera: Ceratopogonidae), anthomyiids (Diptera: Anthomyiidae) and cantharidin-producing oedemerids (Coleoptera: Oedemeridae). - *J. Chem. Ecol.* **20** (8): 1795-1812.
- FRENZEL, M.; HAVELKA, P. & BRANDL, R. 1998: Morphological comparison of canthariphilic Atrichopogon (Diptera: Ceratopogonidae) from Europe and Central Africa - Implications for the evolution of canthariphily? *Zoology* **101**, Supplement I: 37.
- GÖRNITZ, K. 1937: Cantharidin als Gift und Anlockungsmittel für Insekten. - *Arb. phys. angew. Ent.* Berlin-Dahlem **4**: 116-157.
- HEMP, C. 1994: Anthiciden und Cantharidin. Ein Beitrag zur chemischen Ökologie, Bionomie und Phylogenie der Ameisenkäfer (Coleoptera: Anthicidae). Dissertation, Universität Bayreuth. 131 S.
- HEMP, C. & DETTNER, K. 1997: Morphology and Chemistry of Mesothoracic Glands in Anthicid Beetles (Coleoptera: Anthicidae). - *Entomol. Gener.* **22** (2): 97-108.
- HEMP, C. & DETTNER, K. (in press): Attraction of Miridae and Lygaeidae (Heteroptera) to cantharidin. - Ecotropica.
- HEMP, C.; DETTNER, K.; UHMANN, G. & HEMP, A. 1997: A contribution to the biology of the African canthariphilous anthicids *Formicomus rubricollis* LAFERTÉ and *F. gestroi* PIC (Coleoptera: Anthicidae). - *Mitt. Münch. Ent. Ges.* **87**: 81-96.
- HEMP, C.; HEMP, A. & DETTNER, K. 1999a: Canthariphilous Insects in East Africa. - *J. East Afr. Soc.* **88** (1 / 2). In press.
- HEMP, C.; HEMP, A. & DETTNER, K. 1999b: Attraction of the Colour beetle Species *Pallenothriocera rufimembris* by Cantharidin (Coleoptera: Cleridae). - *Entomol. Gen.* **24** (2): 115-123.
- HOLZ, C.; STREIL, G.; DETTNER, K.; DÜTEMEYER, J. & BOLAND, W. 1994: Intersexual 3 of a Toxic Terpenoid during Copulation and its Paternal Allocation to Developmental Stages: Quantification of Cantharidin in Cantharidin-Producing Oedemerids (Coleoptera: Oedemeridae) and Canthariphilous Pyrochroids (Coleoptera: Pyrochroidae). - *Z. Naturforsch.* **49c**: 856-864.
- JUANJIE, T.; YOUWEI, Z.; SHUYONG, W.; ZENGJI, D. & CHUANXIAN, Z. 1995: Investigation on the natural resources and utilization of the Chinese medicinal beetles - Meloidae. - *Acta Entomol. Sinica* **38**: 324-331.

- MAFRÀ-NETO, A. & JOLIVET, P. 1994: Entomophagy in Chrysomelidae: adult *Aristobrotica angulicollis* (ERICHSON) feeding on adult meloids (Coleoptera). - In: Novel Aspects of the Biology of Chrysomelidae (JOLIVET, P. H.; COX, M. L. and PETITPIERRE, E. (eds.)). Kluwer Academic Publishers, Dordrecht, The Netherlands: 171-178.
- MENZEL, F. & MOHRIG, W. 1997: 2.6. Family Sciaridae: 51-69. - In: Contributions to a manual of Palearctic Diptera. Vol. 2. Nematocera and lower Brachycera. (L. PAPP & B. DARVAS (eds.)). Science Herald, Budapest.
- NICHOLLS, D. S. H.; CHRISTMAS, T. I. & GREIG, D. E. 1990: Oedemerid blister beetle dermatosis: A review. - J. Am. Acad. Dermatol. **22**: 815-819.
- PINTO, J. D. 1978: The parasitisation of blister beetles by species of Miridae (Coleoptera: Meloidae; Hemiptera: Miridae). - The Pan Pacific Entomologist **54**: 57-60.
- SCHÜTZ, C. & DETTNER, K. 1992: Cantharidin-Secretion by Elytral Notches of Male Anthicid-Species (Coleoptera: Anthicidae). - Z. Naturforsch. **47c**: 290-299.
- VAN HILLE, J. C. 1954: Cantharidin and Anthicidae. - South African Journal of Science **51**: 154-155.
- VAN HILLE, J. C. 1971: Anthicidae (Coleoptera) from Northern Zululand. - Trans. Roy. Soc. S. Afr. **39** (4): 367-391.
- VAN HILLE, J. C. 1984a: Monograph of *Aulacoderus* LAFERTÉ, a subgenus of *Anthicus* PAYKULL (Coleoptera: Anthicidae). - Annals of the Cape Provincial Museums **15** (1): 1-171.
- VAN HILLE, J. C. 1984b: New south African species of *Anthicus* (*Aulacoderus* LAF.) and *Notoxus* GEOFFR. (Coleoptera: Anthicidae). - Durban Museum Novitates **8** (13): 155-166.
- VAN HILLE, J. C. 1985: Anthicidae (Coleoptera Heteromera) collected in Botswana, 1982-83. - Botswana Notes and Records **17**: 149-162.
- VAN HILLE, J. C. 1988: Three new South African species of *Aulacoderus* LAFERTÉ, a subgenus of *Anthicus* PAYKULL (Coleoptera: Anthicidae). - Annals of the Cape Provincial Museums, Natural History **16** (12): 321-326.
- WIRTH, W. W. 1956: The biting midges ectoparasitic on blister beetles. - Proc. Ent. Soc. Wash. **58**: 15-23.
- WIRTH, W. W. 1980: A new species and corrections in the Atrichopogon midges of the subgenus Melochelea attacking blister beetles (Diptera: Ceratopogonidae). - Proc. Entomol. Soc. Wash. **82**: 124-139.
- YOUNG, D. K. 1981: Bionomics, phylogeny and systematics of the North American species of *Pedilus* FISCHER (Coleoptera: Pyrochroidae). - Thesis, Michigan State University, Dep. of Entomology.
- YOUNG, D. K. 1984a: Cantharidin and insects: an historical review. - The Great Lakes Entomologist **17** (4): 187-194.
- YOUNG, D. K. 1984b: Field records and observations of insects associated with cantharidin. - The Great Lakes Entomologist **17** (4): 195-199.
- YOUNG, D. K. 1989: Notes on the Bionomics of *Xenomyces morrisoni* Horn (Coleoptera: Endomychidae), Another Cantharidin-orienting Fungus Beetle. - Pan-Pacific Entomologist **65** (4): 447-448.

Authors' addresses:

Dr. C. HEMP and Prof. K. DETTNER
 Department of Animal Ecology II
 Bayreuth University
 D - 95440 Bayreuth