

Beitr. Ent.	Keltern	ISSN 0005 - 805X
61 (2011) 1	S. 223 - 238	16.05.2011

# Characterization of leaf-rollers attacking forest and fruit trees in Azerbaijan

## (Lepidoptera: Tortricidae)

With 2 tables, 1 map and 1 figure

SHEYDA MAHARRAMOVA

### Summary

Characterization of the leaf-roller moths (Tortricidae) attacking forest and fruit trees in the North (Sheki-Zagatala), East (Baku-Absheron), and South (Lenkoran-Astara) regions of Azerbaijan was conducted over the decade 1994-2004. Sixteen species of leaf-rollers representing 14 genera were found to be widespread throughout the regions - 9 species of Tortricinae and 7 species of Olethreutinae. Four species, *Psycholoma lecheana* (L.), *Cacoecimorpha pronubana* HÜBNER, *Eudemis profundana* (DENIS & SCHIFFERMÜLLER), and *Epinotia demarniana* (FISCH.), are newly recorded for Azerbaijan. *Tortrix viridana* L. represented 46.0 % of the total number of specimens, *Archips rosanus* L. 28.0 %, and *Archips xylosteanus* L. 20.2 %.

### Zusammenfassung

In der Dekade 1994-2004 wurden im Norden (Sheki-Zagatala), Osten (Baku-Absheron) und Süden (Lenkoran-Astara) Aserbaidschans an Forst- und Obstbäumen schädigende Wickler untersucht. Dabei zeigte sich, dass 16 Wicklerarten aus 14 Gattungen (9 Tortricinae, 7 Olethreutinae) in der Region weit verbreitet sind. Vier weitere Arten, *Psycholoma lecheana* (L.), *Cacoecimorpha pronubana* HÜBNER, *Eudemis profundana* (DENIS & SCHIFFERMÜLLER) und *Epinotia demarniana* (FISCH.), wurden erstmals in Aserbaidschan gefunden. In den Proben hatten Exemplare von *Tortrix viridana* L. einen Anteil von 46.0 %, *Archips rosanus* L. von 28.0 % und *Archips xylosteanus* L. von 20.2 %.

### Key words

Food plants, pest species, leaf-roller, caterpillar, pupa, imago.

### Introduction

Tortricidae are one of the largest families of Microlepidoptera, with more than 9,000 described species (BROWN 2005) distributed worldwide. Information on the tortricids of Azerbaijan has been presented by BOGACHEV (1951), AKHUNDOVA-TUAYEVA (1960), ABDULLAYEVA (1988, 1990), and MAHARRAMOVA (2003, 2006, 2008), with over 200 species of leaf-rollers now recorded from the country (ABDULLAYEVA, 1990).

Leaf rollers are primarily polyphagous and oligophagous. The common name "leaf-rollers" is derived from the fact that the larvae of many species roll the edge of the foliage on which they feed, creating a shelter that may protect them from predators such as birds. This leaf-rolling habit causes serious damage to a variety of forest and fruit trees and ornamental and other plants. Many pest species can be identified by the shape and configuration of the leaf shelter.

The aim of the paper is to present an analysis of the species composition, host associations and yearly distribution of the leaf-rollers in several regions of Azerbaijan. Our object is to determine the host plants of Tortricidae. So we have not included the Tortricidae with unknown host plants.

Characterization of the leaf-roller moths (Tortricidae) attacking forest and fruit trees in the North (Sheki-Zagatala), East (Baku-Absheron), and South (Lenkoran-Astara) regions of Azerbaijan was conducted in 1994-2004. Sixteen species of leaf rollers representing 14 genera were found to be widespread throughout the regions - 9 species of Tortricinae and 7 species of Olethreutinae. *Tortrix viridana* L. represented 46.0 % of the total specimens, *Archips rosanus* L. 28.0 %, and *Archips xylosteanus* L. 20.2 %; *Archips crataeganus* (Hüb), *Ptycholoma lecheana* (L.) and *Aleimma loeflingiana* (L.) each represented less than 1 % of the specimens collected. The purpose of this paper is to detail the results of this survey.

## Material and methods

Field work was conducted from 1994 to 2004 from April to September in the North (Sheki-Zagatala - 1995), East (Baku-Absheron - 1995-1997, 1999-2000, 2004) and South (Lenkoran - 1994-1997, 2001, 2003-2004; Astara - 1994-1998; Masalli - 1995-1996, 1998) regions of Azerbaijan. Forests, orchards, gardens, and roadsides were investigated within the research areas; Hirkan National Park also was sampled. The paper includes the leaf-rollers found on the host plants (trees and bushes only) in larval or pupal stage. Collected larvae and pupae were kept in the laboratory until imagines hatched. Species determination was done according to these imagines. Species were collected in nature in the larval and pupal stage.



Fig. 1: Map of Azerbaijan: o designations; ▲ - material collected area.

Each larva picked from the host plant was kept in a separate jar in the laboratory and fed on host plant leaves. Development of each specimen was observed up to imago stage (FASULATI, 1971). Laboratory work was conducted at the Institute of Zoology of the National Academy of Sciences of Azerbaijan. All specimens are deposited in that institution.

### Study sites:

Forests in the Sheki-Zagatala region of Azerbaijan (Fig. 1) are characterized by a predominance of fruit trees, forest trees, and shrubs. The Baku and Absheron peninsulas in eastern Azerbaijan are characterized by a variety of endemic plants but also by invasives introduced from different countries and other regions of Azerbaijan. The southern regions of Azerbaijan (Lenkoran, Astara, and Masalli) are characterized by 36 species of endemic and relic plants of the tertiary era and 40 species of plants dating from of quaternary period.

## Results

### Subfamily Tortricinae

#### *Tribe Archipini*

#### **Genus *Pandemis* HÜBNER, 1825**

#### ***Pandemis cerasana* (HÜBNER, 1786)**

#### Specimens examined:

**Azerbaijan:** Astara District: Tengerud vil., 19 May 1995, *Malus* sp., 2 ♀♀; Tengerud vil., 19 May 1995, *Malus* sp., 2 ♂♂; Serek vil., 19 May 1995, *Malus* sp., 1 ♀; Serek vil., 20 May 1995, *Malus* sp., 1 ♂; Serek vil., 24 May 1995, *Crataegus* sp., 1 ♀; Astara, 25 May 1995, *Crataegus* sp., 1 ♀.

Larvae and pupae of this species were discovered in the field. The larvae damage leaves and fruit near leaves of the host; the primary host is *Malus* sp. (Table 2). Caterpillars fold the edge of the leaf and bend it toward the mid-vein or spin leaves together with silk. Caterpillars skeletonize leaves and occasionally damage the fruit pedicel which causes the fruit to decay. Caterpillars pupate in the same place as they feed. Females lay eggs in groups of 20-100 on the upper side of the leaf.

#### **Genus *Archips* HÜBNER, 1822**

#### ***Archips rosanus* (LINNAEUS, 1758)**

#### Specimens examined:

**Azerbaijan:** Lenkeran District, 20 September 1994, *Quercus castaneifolia*, 1 ♀. Sheki District, Dodu v., 22 April 1995, *Malus* sp., 10 ♀♀, 5 ♂♂; Sheki District, 22 April 1995, *Malus* sp., 1 ♂; 23 April 1995, *Prunus domestica*, 1 ♀; 11 May 1995, *Armeniaca vulgaris*, 1 ♀; Astara District, Archivan v., 14 May 1995, *Cydonia* sp., 2 ♀♀; Astara District, 15 May 1995, *Malus* sp., 1 ♂; Sheki District, Dodu v., 16 May 1995, *Malus* sp., 1 ♂; Astara District, Archivan v., 17 May 1995, *Cydonia* sp., 3 ♂♂; 20 May 1995, *Cydonia* sp., 1 ♀; 20 May 1995, *Malus* sp., 1 ♀; Sheki District, 21 May 1995, *Malus* sp., 1 ♂; Astara District, Serek v., 24 May 1995, *Crataegus* sp., 3 ♀♀, 5 ♂♂; Sheki District, 24 May 1995, *A. vulgaris*, 2 ♀♀; Baku city, 24 May 1995, *A. vulgaris*, 9 ♀♀; 24 May 1995, *Malus* sp., 1 ♂; 29 May 1995, *Malus* sp., 3 ♀♀, 10 ♂♂; Absheron penin., Merdekan sett.,

14 May 1996, *Cydonia* sp., 1 ♀; Baku city, 16 May 1996, *A. vulgaris*, 12 ♀ ♀, 10 ♂ ♂; 30 July 1996, *Cydonia* sp., 1 ♂; 13 August 1996, *Ulmus suberosa*, 1 ♂; Astara District, Arcivan v., 14 May 1997, *Q. castaneifolia*, 1 ♀; 14 May 1997, *Malus* sp., 2 ♀ ♀; Baku city, 10 June 1997, *Amorpha fruticosa*, 10 ♀ ♀, 7 ♂ ♂; 10 June 1997, *Tilia* sp., 3 ♂ ♂; 10 June 1997, *Fraxinus* sp., 1 ♀; 10 June 1997, *Colutea* sp., 1 ♀; Astara District, Serek v., 10 June 1997, *Q. castaneifolia*, 1 ♀; Baku city: 11 June 1997, *Maclura aurantica*, 1 ♀; 11 June 1997, *Crataegus* sp., 1 ♂; 12 June 1997, *Colutea* sp., 1 ♀; 12 June 1997, *Tilia* sp., 1 ♂; 12 June 1997, *Ligustrum* sp., 6 ♀ ♀; 17 June 1997, *A. fruticosa*, 1 ♀; 18 June 1997, *U. suberosa*, 1 ♂; 23 June 1997, *Punica granatum*, 1 ♀; 15 June 1999, *A. vulgaris*, 1 ♀; Lenkeran District, Hirkan National Park: 7 May 2001, *P. persica*, 1 ♂; 7 May 2001, *Alnus barbata*, 1 ♀; Lenkeran District, Khanbulan area, 10 May 2001, *Salix kusnetzowii*, 1 ♀; 10 May 2001, *Rubus* sp., 2 ♂ ♂; 11 May 2001, *Rubus* sp., 2 ♂ ♂; 11 May 2001, *S. kusnetzowii*, 2 ♀ ♀; Lenkeran District, Hirkan National Park, 15 May 2001, *Gleditsia caspia*, 2 ♂ ♂; Lenkeran District, Khanbulan area, 15 May 2001, *S. kusnetzowii*, 2 ♀ ♀; Lenkeran District, Hirkan National Park: 15 May 2001, *A. barbata*, 1 ♂; 15 May 2001, *Q. castaneifolia*, 1 ♀; Lenkeran District, Khanbulan area, 16 May 2001, *Q. castaneifolia*, 2 ♀ ♀; 16 May 2001, *S. kusnetzowii*, 2 ♀ ♀, 1 ♂; Lenkeran District, Hirkan National Park, 16 May 2001, *Parrotia persica*, 1 ♂; 16 May 2001, *Carpinus caucasica*, 1 ♂; Lenkeran District, Khanbulan area: 16 May 2001, *Rubus* sp., 1 ♀; 17 May 2001, *P. persica*, 1 ♀; 17 May 2001, *Rubus* sp., 1 ♂; Lenkeran District, Hirkan National Park, 18 May 2001, *Q. castaneifolia*, 7 ♂ ♂, 7 ♀ ♀; 18 May 2001, *Zelkova carpinifolia*, 1 ♂; Lenkeran District, Khanbulan area, 18 May 2001, *Rubus* sp., 1 ♀; Lenkeran District, Hirkan National Park, 21 May 2001, *P. persica*, 1 ♂; 23 May 2001, *Z. carpinifolia*, 1 ♀; 24 May 2001, *Q. castaneifolia*, 10 ♀ ♀; 29 May 2001, *P. persica*, 1 ♀; 29 May 2001, *Q. castaneifolia*, 2 ♀ ♀; 29 May 2001, *A. barbata*, 1 ♂; 29 May 2001, *A. barbata*, 1 ♀; Lenkeran District, Hirkan National Park, Druzavel area, 29 May 2001, *Robina pseudoacacia*, 1 ♀; Lenkeran District, Hirkan National Park, 29 May 2001, *Q. castaneifolia*, 1 ♀; 29 May 2001, *S. kusnetzowii*, 2 ♂ ♂; 29 May 2001, *Q. castaneifolia*, 1 ♀; 29 May 2001, *Gleditsia caspia*, 2 ♀ ♀; 29 May 2001, *Mespilus* sp., 2 ♂ ♂; Lenkeran District, Xanbulan area, 29 May 2001, *Acer hyrcanum*, 1 ♀; Lenkeran District, Hirkan National Park, Druzavel area, 29 May 2001, *Robina pseudoacacia*, 1 ♀, 1 ♂; Lenkeran District, Hirkan National Park, 30 May 2001, *Q. castaneifolia*, 3 ♀ ♀, 1 ♂; Lenkeran District, Khanbulan area, 30 May 2001, *Q. castaneifolia*, 1 ♀ 1 ♂; Lenkeran District, Hirkan National Park: 30 May 2001, *Q. castaneifolia*, 3 ♂ ♂; 30 May 2001, *C. caucasica*, 2 ♀ ♀; 1 June 2001, *P. persica*, 4 ♀ ♀; 01 June 2001, *A. barbata*, 3 ♂ ♂; Lenkeran District, Khanbulan area, 1 June 2001, *P. persica*, 2 ♀ ♀, 3 ♂ ♂; 1 June 2001, *P. persica*, 1 ♂; 1 June 2001, *Q. castaneifolia*, 4 ♀ ♀, 1 ♂; Lenkeran District, Hirkan National Park: 5 June 2001, *Q. castaneifolia*, 1 ♂; 5 June 2001, *P. persica*, 1 ♀; 05 June 2001, *Z. carpinifolia*, 1 ♀; 5 June 2001, *P. persica*, 1 ♂; 5 June 2001, *Q. castaneifolia*, 1 ♀; 29 April 2002, *Z. carpinifolia*, 1 ♂; 25 May 2002, *P. persica*, 1 ♂; Absheron penins., Merdekan sett., 24 June 2003, *Q. castaneifolia*, 1 ♀; Lenkeran District, Azfilial area, 24 May 2004, *Planatus digitifolia*, 2 ♀ ♀; Lenkeran District, Hirkan National Park, 25 May 2004, *Ligustrum* sp., 2 ♀ ♀; Baku city, 31 May 2004, *Fraxinus* sp., 1 ♀, 1 ♂; Lenkeran District, 31 May 2004, *Q. castaneifolia*, 2 ♀ ♀, 1 ♂; 31 May 2004, *P. persica*, 1 ♀, 1 ♂; Lenkeran District, Khanbulan area, 1 June 2004, *Rubus* sp., 10 ♀ ♀, 1 ♂; Baku city, 7 June 2004, *Fraxinus* sp., 3 ♀ ♀; 8 June 2004, *Fraxinus* sp., 1 ♂.

We discovered caterpillars and pupae in the field. Larvae of *A. rosanus* roll one to several leaves and join them by silk. When the shelter includes one leaf, it is rolled from the edge towards the mid-vein. Larvae feed on the parenchyma, leaving the veins. *A. rosanus* damages primarily terminal buds and shoots, but larvae attack buds, flowers, leaves, and fruits. Twenty-six plant species were recorded as hosts, including all deciduous trees, forest plants, ornamental plants, shrubs, subtropical plants, and berries, with *Quercus castaneifolia*, *Malus* sp., and *Armeniaca vulgaris* as the main food plants.

Our observations indicate that *A. rosanus* is univoltine. Each egg mass contains up to 60 eggs; the average productivity of a female is 250 eggs. This species overwinters in the egg stage. Bursting of the apple tree buds serves as a phenological signal for the pest to break winter diapause (MAHARRAMOVA, 2003).

### *Archips crataeganus* (HÜBNER, [1799])

#### Specimens examined:

**Azerbaijan:** Lenkeran District, 1 June 2001, *Rubus* sp., 1 ♀.

Rather than rolling the leaves of the host, *Archips crataeganus* joins the axial leaves of a branch with silk creating a nest within which it feeds.

### *Archips xylosteanus* (LINNAEUS, 1758)

#### Specimens examined:

**Azerbaijan:** Sheki District, Dodu v., 22 April 1995, *Malus* sp., 1 ♀; Baku city, 16 May 1996, *Armeniaca vulgaris*, 1 ♂; Lenkeran District, Hirkan National Park, 7 May 2001, *Q. castaneifolia*, 1 ♀, 1 ♂; Lenkeran District, Khanbulan area, 8 May 2001, *Alnus subcordata*, 1 ♀; Lenkeran District, Hirkan National Park, 8 May 2001, *Populus hyrcana*, 2 ♂♂; 10 May 2001, *P. hyrcana*, 1 ♂; 11 May 2001, *P. hircana*, 1 ♂; 11 May 2001, *Carpinus caucasica*, 1 ♀; 11 May 2001, *Q. castaneifolia*, 1 ♀; 15 May 2001, *Q. castaneifolia*, 1 ♀; 15 May 2001, *C. caucasica*, 2 ♂♂; 15 May 2001, *P. hircana*, 1 ♀; Lenkeran District, Khanbulan, 15 May 2001, *Q. castaneifolia*, 1 ♀; Lenkeran District, Hirkan National Park: 16 May 2001, *Parrotia persica*, 1 ♀; 16 May 2001, *C. caucasica*, 1 ♀; Lenkeran District, Khanbulan area, 16 May 2001, *Q. castaneifolia*, 2 ♂♂; 16 May 2001, *P. persica*, 2 ♀♀; Lenkeran District, Hirkan National Park, 16 May 2001, *Mespilus* sp., 1 ♀; Lenkeran District, Khanbulan area, 17 May 2001, *P. persica*, 1 ♂; Lenkeran District, Hirkan National Park: 18 May 2001, *Z. carpinifolia*, 5 ♀♀, 1 ♂; 18 May 2001, *Q. castaneifolia*, 14 ♀♀; 18 May 2001, *P. persica*, 1 ♀; 29 May 2001, *Mespilus* sp., 3 ♂♂; 29 May 2001, *Populus hyrcana*, 1 ♀; 29 May 2001, *Z. carpinifolia*, 5 ♀♀; 29 May 2001, *Q. castaneifolia*, 3 ♀♀; 29 May 2001, *C. caucasica*, 3 ♂♂; Lenkeran District, Khanbulan area: 29 May 2001, *P. persica*, 1 ♀; 30 May 2001, *Q. castaneifolia*, 9 ♀♀, 9 ♂♂; 30 May 2001, *C. caucasica*, 3 ♀♀; 30 May 2001, *Q. castaneifolia*, 2 ♀♀, 3 ♂♂; Lenkeran District, Hirkan National Park, 1 June 2001, *Q. castaneifolia*, 31 ♀♀, 11 ♂♂; Lenkeran District, Khanbulan area, 1 June 2001, *Q. castaneifolia*, 1 ♀; Lenkeran District, Hirkan National Park, 1 June 2001, *P. persica*, 7 ♂♂; 1 June 2001, *C. caucasica*, 3 ♀♀; 5 June 2001, *Populus hyrcana*, 3 ♂♂; 5 June 2001, *P. persica*, 1 ♀; 27 April 2002, *Q. castaneifolia*, 3 ♀♀; 29 April 2002, *P. persica*, 10 ♀♀, 3 ♂♂; 29 April 2002, *Q. castaneifolia*, 13 ♀♀, 4 ♂♂; 29 April 2002, *Z. carpinifolia*, 2 ♀♀; 1 May 2002, *Q. castaneifolia*, 1 ♂; 3 May 2004, *Q. castaneifolia*, 3 ♂♂; 25 May 2004, *Q. castaneifolia*, 1 ♀; Lenkeran District, Khanbulan area, 31 May 2004, *Q. castaneifolia*, 4 ♂♂; 1 June 2004, *Rubus* sp., 1 ♀.

Caterpillars and pupae of *Archips xylosteanus* were collected in nature. This species is a defoliator of a wide range of broadleaf trees. Larvae are omnivorous and are known to damage orchard, ornamental, and forest trees. Females lay eggs in groups of 2-100 eggs on the trunk of the tree or on branches near the ground. The phenological signal for caterpillar hatching is opening of the apple tree buds. Caterpillars roll the leaf beginning from the top of the leaf along its width. We documented larval damage on more than 10 species of plants, but especially *Q. castaneifolia*, *P. persica*, and *Z. carpinifolia* (Table 2). *A. xylosteanus* was recorded throughout the study area and survey period; however, it was most numerous in 2001-2002 in Lenkeran.

**Genus *Ptycholoma* STEPHENS, 1929*****Ptycholoma lecheana* (LINNAEUS, 1758)****Specimens examined:**

**Azerbaijan:** Absheron penins., Merdekan settl., 22 May 2001, *Q. castaneifolia*, 1 ♀.

Early instars skeletonize leaves; later instars damage buds, flowers, and newly emerged leaves. Larvae join several new leaves by silk and feed inside.

**Genus *Cacoecimorpha* OBRAZTSOV, 1954*****Cacoecimorpha pronubana* HÜBNER, 1800****Specimens examined:**

**Azerbaijan:** Absheron penin., Merdekan settl., 6 June 1997, *Laurus* sp., 1 ♀; Lenkeran District, Hirkan National Park, 27 June 2003, *Ligustrum* sp., 1 ♂.

Larvae were collected in nature. Caterpillars spin together 2-3 leaves with silk and eat the parenchyma tissue inside.

**Tribe Tortricini****Genus *Aleimma* HÜBNER, [1825]*****Aleimma loeflingiana* (LINNAEUS, 1758)****Specimens examined:**

**Azerbaijan:** Lenkeran District, 24 June 2003, *Q. castaneifolia*, 1 ♀.

The caterpillar wraps leaves from the main rib to the lower face obliquely and ties them with silk.

**Genus *Tortrix* LINNAEUS, 1758*****Tortrix viridana* LINNAEUS, 1758****Specimens examined:**

**Azerbaijan:** Lenkeran District, 20 April 1994, *Q. castaneifolia*, 1 ♀; Lenkeran District, Hirkan National Park: 30 April 1994, *Q. castaneifolia*, 1 ♀; 5 May 1994, *Q. castaneifolia*, 1 ♀; Lenkeran District, Khanbulan area, 20 April 1994, *Q. castaneifolia*, 1 ♀, 2 ♂♂; Lenkeran District: 10 May 1995, *Zelkova carpinifolia*, 2 ♂♂; 10 May 1995, *Q. castaneifolia*, 1 ♂; Sheki District, Markhal area, *Quercus* sp., 15 May 1995, 1 ♀; Astara District: 15 May 1995, *Q. castaneifolia*, 1 ♀; 17 May 1995, *Q. castaneifolia*, 1 ♂; Astara District, *Q. castaneifolia*, Serek v., 18 May 1995, Astara District, *Q. castaneifolia*, 1 ♂; Astara District, Serek v., 19 May 1995, *Ulmus suberosa*, 1 ♀; 19 May 1995, 10 ♀♀, 2 ♂♂; 19 May 1995, *Carpinus caucasica*, 1 ♀; 19 May 1995, *P. persica*, 1 ♂; Astara District, Tengerud v., 19 May 1995, *Q. castaneifolia*, 9 ♀♀; 19 May 1995, *Tilia* sp., 3 ♂♂; Astara District, 19 May 1995, *Q. castaneifolia*, 2 ♂♂; Masally District, 20 May 1995, *Q. castaneifolia*, 20 ♀♀, 6 ♂♂; 21 May 1995, *Q. castaneifolia*, 3 ♀♀, 1 ♂♂; Sheki District, Markhal area, 24 May 1995, *Quercus* sp., 1 ♂; Masally District, 24 May 1995, *Q. castaneifolia*, 1 ♂; 27 May 1995, *Q. castaneifolia*, 5 ♀♀; 28 May 1995, *Q. castaneifolia*, 17 ♀♀, 5 ♂♂; 24 April 1996, *Q. castaneifolia*, 1 ♀; Lenkeran District, Hirkan National Park: 2 May 1996,

*Q. castaneifolia*, 1 ♂; 5 May 1996, *Q. castaneifolia*, 1 ♂; Lenkeran District: 15 May 1996, *Q. castaneifolia*, 1 ♀; 15 June 1996, *Q. castaneifolia*, 1 ♀; Astara District, Archivan v., 14 May 1997, *Q. castaneifolia*, 6 ♀ ♀, 2 ♂ ♂; Astara District, 16 May 1997, *Q. castaneifolia*, 1 ♀; Astara District, Tengerud v., 19 May 1997, *Q. castaneifolia*, 1 ♂; Baku city, 18 June 1997, *Ulmus suberosa*, 1 ♂; Lenkeran District: 22 June 1997, *Q. castaneifolia*, 1 ♂; 27 June 1997, *Z. carpinifolia*, 1 ♀; Masally District, 22 May 1998, *Q. castaneifolia*, 1 ♀; Lenkeran District, Hirkan National Park: 29 April 2001, *Populus hyrcana*, 1 ♀; 02 May 2001, *Q. castaneifolia*, 1 ♂; 3 May 2001, *Zelkova carpinifolia*, 1 ♂; 4 May 2001, *Cydonia* sp., 1 ♀; 4 May 2001, *Q. castaneifolia*, 2 ♀ ♀, 3 ♂ ♂; 4 May 2001, *Z. carpinifolia*, 5 ♂ ♂; 4 May 2001, *Populus hyrcana*, 1 ♀; 4 May 2001, *Mespilus* sp., 1 ♂, 1 ♀; 5 May 2001, *Parrotia persica*, 1 ♂; 5 May 2001, *Mespilus* sp., 1 ♀; 6 May 2001, *Cydonia* sp., 1 ♀; 6 May 2001, *Q. castaneifolia*, 4 ♀ ♀, 6 ♂ ♂; 6 May 2001, *Z. carpinifolia*, 5 ♀ ♀, 2 ♂ ♂; 6 May 2001, *P. hyrcana*, 4 ♀ ♀; 6 May 2001, *Alnus barbata*, 3 ♂ ♂; 6 May 2001, *Mespilus* sp., 2 ♀ ♀, 1 ♂; 6 May 2001, *Crataegus* sp., 4 ♀ ♀, 7 May 2001, *P. persica*, 1 ♂; 7 May 2001, *Q. castaneifolia*, 10 ♀ ♀, 2 ♂ ♂; 8 May 2001, *Q. castaneifolia*, 6 ♀ ♀, 20 ♂ ♂; 8 May 2001, *U. suberosa*, 10 ♀ ♀, 21 ♂ ♂; 8 May 2001, *Carpinus caucasica*, 2 ♀ ♀, 3 ♂ ♂; 8 May 2001, *Parrotia persica*, 5 ♀ ♀, 4 ♂ ♂; 8 May 2001, *Cydonia* sp., 1 ♀, 1 ♂; 8 May 2001, *Mespilus* sp., 1 ♀; 9 May 2001, *Q. castaneifolia*, 2 ♀ ♀, 2 ♂ ♂; Lenkeran District, Khanbulan area, 9 May 2001, *P. persica*, 1 ♀; Lenkeran District, Hirkan National Park, 9 May 2001, *Z. carpinifolia*, 2 ♂ ♂; 9 May 2001, *Mespilus* sp., 1 ♀; 10 May 2001, *Zelkova carpinifolia*, 1 ♀; 10 May 2001, *Q. castaneifolia*, 4 ♀ ♀; 10 May 2001, *P. persica*, 3 ♂ ♂; 10 May 2001, *Z. carpinifolia*, 2 ♀ ♀; Lenkeran District, Khanbulan area, 10 May 2001, *P. persica*, 2 ♂ ♂; Lenkeran District, Hirkan National Park: 10 May 2001, *Crataegus* sp., 1 ♀; 11 May 2001, *Z. carpinifolia*, 2 ♀ ♀; 11 May 2001, *Mespilus* sp., 1 ♂; 11 May 2001, *Ulmus suberosa*, 1 ♀; 11 May 2001, *Q. castaneifolia*, 2 ♀ ♀; 14 May 2001, *Q. castaneifolia*, 1 ♂; 15 May 2001, *P. persica*, 2 ♀ ♀, 1 ♂; 15 May 2001, *Q. castaneifolia*, 1 ♀; 15 May 2001, *U. suberosa*, 1 ♀; 15 May 2001, *Cydonia* sp., 1 ♂; Lenkeran District, Khanbulan area, 15 May 2001, *P. persica*, 2 ♀ ♀; Lenkeran District, Hirkan National Park, 16 May 2001, *Cydonia* sp., 1 ♂; Lenkeran District, Khanbulan area, 16 May 2001, *P. persica*, 2 ♀ ♀; 16 May 2001, *Q. castaneifolia*, 2 ♂ ♂; Lenkeran District, Hirkan National Park: 16 May 2001, *Z. carpinifolia*, 1 ♀, 1 ♂; 16 May 2001, *Mespilus* sp., 1 ♀; 16 May 2001, *Q. castaneifolia*, 1 ♂; 17 May 2001, *Crataegus* sp., 1 ♂; 17 May 2001, *Carpinus caucasica*, 1 ♀; 17 May 2001, *P. persica*, 1 ♀; 17 May 2001, *Cydonia* sp., 1 ♂; Lenkeran District, Khanbulan area, 17 May 2001, *P. persica*, 2 ♀ ♀; 18 May 2001, *P. persica*, 1 ♂; Lenkeran District, Hirkan National Park: 18 May 2001, *Carpinus caucasica*, 1 ♂; 18 May 2001, *Crataegus* sp., 1 ♀; 18 May 2001, *Zelkova carpinifolia*, 3 ♀ ♀; 18 May 2001, *Q. castaneifolia*, 9 ♀ ♀, 2 ♂ ♂; 21 May 2001, *P. persica*, 8 ♀ ♀, 1 ♂; 21 May 2001, *Cydonia* sp., 1 ♀, 1 ♂; 21 May 2001, *Ulmus suberosa*, 8 ♀ ♀; 22 May 2001, *Carpinus caucasica*, 4 ♀ ♀; 22 May 2001, *Q. castaneifolia*, 9 ♀ ♀, 2 ♂ ♂; 24 May 2001, *Q. castaneifolia*, 7 ♀ ♀, 3 ♂ ♂; 29 May 2001, *Q. castaneifolia*, 7 ♀ ♀; 29 May 2001, *Crataegus* sp., 1 ♀, 2 ♂ ♂; 29 May 2001, *Mespilus* sp., 1 ♀; 30 May 2001, *Q. castaneifolia*, 10 ♀ ♀, 2 ♂ ♂; 1 June 2001, *Q. castaneifolia*, 10 ♀ ♀, 3 ♂ ♂; Lenkeran District, Khanbulan area, 1 June 2001, *P. persica*, 2 ♀ ♀; Lenkeran District, Hirkan National Park: 1 June 2001, *U. suberosa*, 1 ♀; 5 June 2001, *U. suberosa*, 6 ♀ ♀; 1 May 2002, *Q. castaneifolia*, 1 ♂; 3 May 2004, *Z. carpinifolia*, 2 ♂ ♂; 3 May 2004, *Q. castaneifolia*, 3 ♀ ♀, 4 ♂ ♂; 18 May 2004, *Q. castaneifolia*, 1 ♀, 1 ♂; 19 May 2004, *Q. castaneifolia*, 1 ♂.

We collected caterpillars, pupae, and adults of *Tortrix viridana*. This species overwinters as an egg, which remains dormant until the host leaf buds begin to open in mid-March or early April of the following year. The young caterpillars feed on tender leaves until late April or early May. The fully grown caterpillars eat the leaves entirely. Pupation takes place inside the rolled leaves. Moths begin to emerge about the first of May. Peak flight occurs about 10-15 May. Females lay

up to 40-120 eggs, depending on host plant. Females begin to lay egg mass containing 2 eggs on leaves or twigs of trees completing the one-year life cycle. The literature provides somewhat varying information on host plants of *T. viridana*. Some authors (e.g., LOZOVY, 1965; ABDULLAYEVA, 1990; RAYXOLF-RIM, 2002) considered it to be monophagous, feeding on only one or two species of oak; other authors consider the species to be oligophagous, feeding on several species of oak (ZEROVA et al., 1989). However, according to VORONSOV (1967), the young caterpillars have been recorded from oak, lime, and poplar. We documented *T. viridana* on 12 different tree species in Azerbaijan (Table 2); *Q. castaneifolia*, *Z. carpinifolia*, *U. suberosa*, and *P. persica* are attacked most often by *T. viridana* (MAHARRAMOVA, 2008).

Larval survival of this species is 58 % on *Q. castaneifolia*, 59 % on *Z. carpinifolia*, 82 % on *U. suberosa*. Female productivity varied depending on host. It was 43.5 on *Quercus*, 46 on *P. persica*, 49.6 on *Z. carpinifolia*, 49.6 on *C. caucasica*, and 54 on *U. suberosa*.

### Genus *Croesia* HÜBNER, [1825]

#### *Croesia forskaleana* (LINNAEUS, 1758)

##### Specimens examined:

**Azerbaijan:** Lenkeran District, Khanbulan area, 29 May 2001, *Acer hyrcanum*, 5 ♀♀, 8 ♂♂.

Overwintering caterpillars damage the first flowers of the host and then roll the leaves. Adults fly from May to August. Caterpillars were collected in nature.

## Subfamily Olethreutinae

### Tribe Olethreutini

#### Genus *Eudemis* HÜBNER, [1825]

#### *Eudemis profundana* ([DENIS & SCHIFFERMÜLLER], 1775)

##### Specimens examined:

**Azerbaijan:** Lenkeran District, Hirkan National Park, 25 May 2003, *Q. castaneifolia*, 3 ♀♀; Lenkeran District, 9 June 2004, *Q. castaneifolia*, 2 ♂♂.

Caterpillars damage *Q. castaneifolia*.

#### Genus *Hedya* HÜBNER, [1825]

#### *Hedya nubiferana* HAWORTH, [1811]

##### Specimens examined:

**Azerbaijan:** Sheki District, Markhal area, 22 April 1995, *Quercus* sp., 1 ♀; Sheki District, 11 May 1995, *Armeniaca vulgaris*, 1 ♀; Baku city, 24 May 1995, *Malus* sp., 1 ♀; 24 May 1995, *A. vulgaris*, 2 ♂♂; Sheki city, 28 May 1995, 2 ♀♀; Baku city, 29 May 1995, *Prunus domestica*, 1 ♂; Astara District, Tengerud v., 5 May 1997, *Cydonia* sp., 1 ♀; Astara District, Archivan v., 7 May 1998, *Malus* sp., 1 ♂; Astara District, Serek v., 14 May 1998, *Malus* sp., 1 ♀.

Caterpillars and pupae were collected in nature. This species damages trees and bushes in the family Rosaceae. Of the five recorded hosts, *Malus* sp. and *A. vulgaris* were preferred (Table 1).



This species overwinters as an early instar larva. Overwintering caterpillars hatch in the second week of April and feed on buds; they later attack leaves. Pupation occurs at the end of April and early May. Females lay about 150 eggs on leaves or occasionally on fruit, either or in groups of 2-8. There is one generation annually.

### **Tribe Eucosmini**

#### **Genus *Spilonota* STEPHENS, 1829**

##### ***Spilonota ocellana* ([DENIS & SCHIFFERMÜLLER], 1775)**

#### **Specimens examined:**

**Azerbaijan:** Absheron penins., Merdekan sett.: 20 May 1995, *Malus* sp., 1 ♀; 30 May 1995, *Malus* sp., 1 ♀; Lenkeran District, Hirkan National Park, 29 April 2002, *Q. castaneifolia*, 1 ♀.

We collected caterpillars in nature. This species overwinters as a caterpillar starting in September-October. Bud opening serves as the phenological signal for caterpillars to emerge in the spring. Early instars damage buds, and later instars feed on young leaves. Pupation takes place within the rolled leaves. Adults appear in June-July.

#### **Genus *Epinotia* HÜBNER, [1825]**

##### ***Epinotia demarniana* (FISCHER VON RÖSLERSTAMM, 1840)**

#### **Specimens examined:**

**Azerbaijan:** Lenkeran District, Hirkan National Park, 1 June 2001, *Alnus barbata*, 1 ♀; Lenkeran District, Hirkan National Park, 1 June 2001, *A. barbata*, 1 ♀.

#### **Genus *Gypsonoma* MYRICK, 1895**

##### ***Gypsonoma* sp.**

#### **Specimens examined:**

**Azerbaijan:** Absheron, Merdekan, 22 June 1996, *Populus* sp., 4 ♀ ♀.

#### **Genus *Notocelia* HÜBNER, [1825]**

##### ***Notocelia uddmanniana* (LINNAEUS, 1758)**

#### **Specimens examined:**

**Azerbaijan:** Lenkeran District, Hirkan National Park, 1 June 2001, *Rubus* sp., 2 ♂ ♂.

Caterpillars feed on raspberry, strawberry, rose, and dog-rose, joining the leaves of the host by silk into a friable ball in which there may be 2-3 caterpillars. Adults appear in June-July.

### Tribe Grapholitini

#### Genus *Cydia* HÜBNER, [1825]

#### *Cydia pomonella* (LINNAEUS, 1758)

#### Specimens examined:

**Azerbaijan:** Astara District, Archivan v., 25 April 1995, *Malus* sp., 1 ♂; Sheki District, Dodu v., 2 May 1995, *Malus* sp., 1 ♀; Baku city, 14 May 1995, *A. vulgaris*, 1 ♂; Astara District, Archivan v., 29 September 1995, *Malus* sp., 1 ♀; Baku city, 30 July 1996, *Malus* sp., 1 ♀; Astara District, Archivan v., 14 May 1997, *Q. castaneifolia*, 1 ♀.

*Cydia pomonella* is oligophagous; we recorded it from three plant species, with *Malus* sp. the primary host (Table 1). There are two generations per year. Females lay eggs on the underside of the leaf. Productivity of the female is 60-120 eggs. Newly hatched caterpillars penetrate into the pulp of the fruit and make a chamberlike cavity. The following instars feed in the seed cavity. The fourth instar completes feeding and leaves the fruit to pupate. Moths are active at dusk.

### Discussion

#### Analysis of the species composition of the leaf rollers

During studies conducted from 1994-2004 we documented 16 species of leaf-rollers belonging to 14 genera. They belong to two subfamilies: Tortricinae (9 species in 7 genera) and Olethreutinae (7 species in 7 genera). *Tortrix viridana* represented 46.0 % of the total specimens,




Fig. 2: Distribution of *Tortrix viridana*, *Archips rosanus* and *Archips xylosteanus* in different regions of Azerbaijan.

*Archips rosanus* L. 28.0 %, and *Archips xylosteanus* L. 20.2 %. Four species were newly recorded for Azerbaijan (Table 1): *Ptycholoma lecheana*, *Cacoecimorpha pronubana*, *Eudemis profundana*, and *Epinotia demarniana*.

Fourteen species of leaf-rollers (87.5 %) were collected in the Lenkoran–Astara, 9 species (56.2 %) in the Baku–Absheron, and 5 species (33.3 %) in the Sheki–Zaqatala. Five species were found in all three regions. The three most abundant species during the survey (i.e., *Tortrix viridana*, *Archips rosanus*, and *Achips xylosteanus*) were most numerous in Lenkoran–Astara (Fig. 2). Figure 2 shows the share of all collected samples in Lenkoran–Astara, Baku–Absheron and Sheki–Zaqatala regions during the studied period. For example, totally 428 samples of *T. viridana* were collected. From them 425 (99,3 %) distributed in Lenkoran–Astara region, 1 (0,2 %) in the Baku–Absheron and 2 (0,5 %) in the Sheki–Zaqatala region.

*Archips crataeganus* (HÜBN.), *Ptychotoma lecheana* (L.), *Aleimma loeflingiana* (L.) were the least common species, each representing about 0.1 % of the total specimens collected.

Two species (*P. lecheana*, *Gypsonoma* sp.) were recorded only in the Baku–Absheron, and 7 species (*E. profundana*, *A. loeflingiana*, *C. forscaleana*, *P. cerasana*, *E. demardiana*, *N. uddmanniana*, *A. crataegana*) only in Lenkoran.

### Host Associations

Species composition varied from region to region, with many leaf-rollers strongly associated with specific plant species.

Research in the Sheki–Zaqatala region was conducted in 1995 where leaf-rollers were recorded primarily on apple, apricot, plum, and oak. Approximately 56 % of damage was to apple tree and 28 % to oak. Other trees suffered from leaf-rollers totally 2–14 %.

Research in the Baku–Absheron peninsular was conducted during 1995–1997, 1999–2001, 2003–2004. Apricot (26.75 %), indigo (14.65 %), apple (12.1 %) suffered serious damage from leaf-rollers. Ash-tree, *Populus* sp., and *U. campestris* suffered 5.73–8.92 % damage. Other plants suffered considerable less, from 0.64–4.46 %.

Research in the Lenkoran region was conducted in 1994–1998, 2001–2004, and thirteen species of leaf-rollers were documented. They were associated with 21 plant species. Damage caused by leaf-rollers was as follows: *Quercus castaneifolia* whose damage constituted 51.4 % of all the damage caused; *P. persica* (11.3 %), *Z. caprinifolia* (7.5 %), *U. suberosa* (6.1 %). Other plants suffered to a lesser degree (0.12 to 3.47 %) (MAHARRAMOVA, 2008).

In all areas studied, three species of leaf-rollers (*A. rosanus*, *A. xylosteanus*, and *T. viridana*) had a wide range of food plants. *A. rosanus* was documented from 26 species of plants, *A. xylosteanus* from 12, and *T. viridana* from 10. *T. viridana* was the most abundant species in 2000–2002 in the Lenkoran region (in the territory of the Hyrcan National Park, as well), causing serious damage to more than 10 plant species. *T. viridana*, a heat-loving, xerophilous species (ZEROVA et al., 1989), was recorded mainly at the flanks and edges of forest causing up to 80 % damage to host plants. During outbreaks it seriously damaged relict and endemic trees in the Hyrcan National Park including *P. persica* and *Z. carpinifolia* which are representatives of the tertiary and the quaternary periods.

### Yearly distribution

Leaf-rollers were most numerous in 2001, with nine species were recorded (Table 1): eight from the Lenkoran–Astara region and one from the Baku–Absheron region. Five of the species from the Lenkoran–Astara region were uncommon (represented by less than 9 individuals): *A. crataegana*, *C. forskalearia*, *Spilonota ocellana*, *E. demardiana*, and *N. uddmanniana*. The lowest year in terms of species richness was 1994 with only two species recorded. Five species were recorded in 1996, with one (i.e., *T. viridana*) from the Lenkoran–Astara region and 4 (*A. rosanus*, *A. xylosteanus*, *Gypsonoma* sp., *Cydia pomonella*) from the Baku–Absheron region. In 1997, 2003 and 2004 only four species were recorded, all of which were recorded in Lenkoran–Astara region, with *A. rosanus* and *T. viridana* also recorded from the Baku–Absheron. In 1998–2002 three species were recorded (one species in Baku–Absheron two in Lenkoran–Astara region); in 1994 two species were recorded (in Lenkoran–Astara region). In 1999 only one species was recorded (in Lenkoran–Astara region). *A. rosanus* and *T. viridana* were documented every year in all studied period.

### Acknowledgments

I am grateful to Dr. SHARGIYA ABDULLAYEVA (Entomological department of the Institute of Zoology of the NAS of Azerbaijan) and Dr. MUSTAFA OZDEMIR (Plant protection and taxonomy museum, Ankara, Turkey) for determination of tortricids, to Dr. ILHAMA KERIMOVA (Entomological department of the Institute of Zoology of the NAS of Azerbaijan) for translation and to Dr. JOHN BROWN (National Museum of Natural History, Washington, DC, USA) for edition of manuscript.

### References

- ABDULLAYEVA, S. Y. 1988: About of the fauna of leaf rollers (Lepidoptera, Tortricidae) in east of Azerbaijan. – Az AS informations, Biol. Res., series, N4: 69-74. (in Azerb.)
- ABDULLAYEVA, S. Y. 1990: About of the fauna of leaf rollers (Lepidoptera, Tortricidae) in east of Azerbaijan. – Az AS informations, Biol. Res., series, N3: 70-73. (in Azerb.)
- AKHUNDOVA-TUAYEVA, L. M. 1960: K izucheniyu cheshuekrilikh vredyashix drevesno-kustarnikovim porodam v Lenkoranskoy zone. – Uch. zap. AGU named S. M. Kirova, N1. (in Russian)
- BOGACHEV, A. V. 1951: Listovertki - Tortricidae. – In book: Jivotniy Mir Azerbaijana, Baku, Pub. AzCCR, 1951, 382-383. (in Russian)
- BROWN, J. W. 2005: World Catalogue of Insects. Volume 5. Tortricidae (Lepidoptera). – Apollo Books, 741 pp. (in English)
- FASULATI, K. K. 1971: Poleviye izucheniye na zemnikh bezpozvonochnikh. – Izd. Visshaya Shkola. Moskva, 423 pp. (in Russian)
- LOZOVY, D. I. 1965: Vredniye nasekomiye parkovikh i lesoparkovikh nasajdeniy Gruzii. – Tbilisi: 272 pp. (in Russian)
- MAHARRAMOVA, S. M. 2003: The role of parasitoid complex in *Archips rosana* L. (Lepidoptera, Tortricidae) number management. – First Congress of Azerbaijan Society of Zoologists. Elm, Baku 2003: pp. 230-236. (in Azerb.)
- MAHARRAMOVA, S. M. 2006: Leaf-rollers (Lepidoptera: Tortricidae) damaging to fruit-wood cultures of the Lenkoran region. – Proceedings of the Azerbaijan Institute of Zoology, Vol. XXVIII: Baku, Elm: pp. 542-551. (in Azerb.)
- MAHARRAMOVA, S. M. 2008: Factor effecting on outbreak of green leaf roller (*Tortrix viridana* L., Tortricidae). – Proceedings of the Azerbaijan society of Zoologists, Volume 1, Baku, Elm: pp. 291-297. (in Azerb.)

RAYXOLF-RIM, X. 2002: Babochki. – Moscow, Astrel. (in Russian)

VORONSOV, A. I. 1967: Lesnaya entomologia. M. – Visshaya shkola. (in Russian)

ZEROVA, M. D.; KOTENKO, A. G.; SEREGINA, L. Y. & TOLKANITS, V. I. 1989: Entomofagi zelenoy dubovoy listovertki i neparnogo shelkopyada yugo-zapada evropeyskoy chasti SSSR. – Kiev: 200 pp. (in Russian)

**Author's address:**

SHEYDA MAHARRAMOVA  
Institute of Zoology  
National Academy of Sciences  
of Azerbaijan Republic, Az1073  
pr. 1128, bl. 504, Baku, Azerbaijan  
sheyda@bakililar.az

**Subject editor:**

Dr. W. NÄSSIG

Tab.1: The species composition and the number of leaf tortricids in Azerbaijan in the years 1994-2004.

Names of species	Lankaran-Astara region										Baku-Absheron region								Shaki-Zaqatala region			Totally															
	Number of individuals [in specimens]										Number of individuals [in specimens]								Number of individuals (in specimens)		Individuals (in specimens)	%															
	1994	1995	1996	1997	1998	1999	2001	2002	2003	2004	total	Percentage contribution in given region	1995	1996	1997	1998	1999	2001	2003	2004			total	Percentage contribution in given region	1995	1996	1997	1998	1999	2001	2003	2004	total	Percentage contribution in given region			
<i>Pandemis cerasana</i> (Hüb.)	8										8	100																						8	0.9		
<i>Archips rosanus</i> (L.)	1	16	4		1	103	2	20	147		20	56.5							1	6	91												22	22	8.5	260	28
<i>Archips crataeganus</i> (Hüb.)						1			1		1	100																						1	0.1		
<i>Archips xylosteanus</i> (L.)						141	36	9	186		9	99																					1	1	0.5	188	20.2
<i>Psycholoma lechearna</i> (L.)*																																		1	0.1		
<i>Cacoecimorpha pronubana</i> Hüb.*											1	50																						2	0.2		
<i>Aleimma loeflingiana</i> (L.)											1	100																						1	0.1		
<i>Tortrix viridana</i> L.	6	93	5	12	1	295	1	12	425		12	99.3																					2	2	0.5	428	46.02
<i>Croesia forskaleana</i> L.						8			8		8	100																						8	0.9		
<i>Eudemis profundana</i> DEN. & SCHIE.*								3	2	5	5	100																						5	0.53		
<i>Hedya nubiferana</i> HAW.				1	2						3	27.3																					4	4	36.36	11	1.2

Names of species	Lankaran-Astara region										Baku-Absheron region								Shaki-Zaqatala region			Totally				
	Number of individuals [in specimens]										Number of individuals [in specimens]								Number of individuals (in specimens)		Indivudals (in specimens)	%				
	1994	1995	1996	1997	1998	1999	2001	2002	2003	2004	total	1995	1996	1997	1998	1999	2001	2003	2004	total			1995	total	Percentage contribution in given region	
<i>Spilonota ocellana</i> DEN. & SCHIE.						1					1	33.3	2						2	66.7				3	0.32	
<i>Epinotia demarniana</i> (FISCH.)*						2					2	100												2	0.2	
<i>Gypsonoma</i> sp.													4						4	100				4	0.43	
<i>Notocelia uddmanniana</i> (L.)						2					2	100												2	0.2	
<i>Cydia pomonella</i> (L.)	2	1										50	1	1					2	33.3			1	1	0.6	
Total	7	119	5	18	3	1	553	39	5	43	793	-	30	31	36	1	1	1	6	107	-		30	30	930	100
Percentage contribution of all species in given site	85.3										11.5								3.2							

\* - New to Azerbaijan fauna

Tab. 2: Food connection of leaf rollers (1994-2004) in Azerbaijan.

Leaf rollers	Leaf rollers recording years	Number of host plants	Host plants & number of leaf roller samples on them	Preferred host plants of leaf rollers
<i>Pandemis cerasana</i>	1995	2	<i>Crataegus</i> sp. - 2, <i>Malus</i> sp. - 6	<i>Malus</i> sp.
<i>Archips rosanus</i>	1994, 1995, 1996, 1997, 1999, 2001, 2002, 2003, 2004	26	<i>Q. castaneifolia</i> - 51, <i>Malus</i> sp. - 36, <i>A. vulgaris</i> - 35, <i>P. persica</i> - 20, <i>Rubus</i> L. - 18, <i>A. fruticosus</i> - 18, <i>Crataegus</i> sp. - 9, <i>Fraxinus</i> sp. - 7, <i>Z. carpinifolia</i> - 4, <i>Mespilus</i> Z. - 2, <i>C. caucasica</i> - 3, <i>A. barbata</i> - 7, <i>A. hyrcanum</i> - 1, <i>R. pseudo-acacia</i> - 3, <i>G. caspia</i> - 4, <i>P. digitifolia</i> - 2, <i>Ligustrum</i> sp. - 8, <i>S. Kusnetzowii</i> - 9, <i>Colutea</i> sp. - 2, <i>Tilia</i> sp. - 4, <i>M. aurantica</i> - 1, <i>P. granatum</i> L. - 1, <i>Cydonia</i> sp. - 8, <i>P. domestica</i> - 1, <i>U. suberosa</i> - 2	<i>Q. castaneifolia</i> <i>Malus</i> sp. <i>A. vulgaris</i> <i>P. persica</i> <i>Rubus</i> L. <i>A. fruticosus</i>
<i>Archips crataegana</i>	2001	1	<i>Rubus</i> sp. - 1	
<i>Archips xylosteanus</i>	1995, 1996, 2001, 2002, 2004	10	<i>Q. castaneifolia</i> - 119, <i>P. persica</i> - 27, <i>C. caucasica</i> - 11, <i>P. hyrcana</i> - 9, <i>Z. carpinifolia</i> - 13, <i>Mespilus</i> Z. - 4, <i>Malus</i> sp. - 1, <i>Rubus</i> L. - 1, <i>A. vulgaris</i> - 1, <i>A. subcordata</i> - 1	<i>Q. castaneifolia</i> <i>P. persica</i> <i>Z. carpinifolia</i> <i>C. caucasica</i>
<i>Ptycholoma lecheana</i>	2001	1	<i>Q. castaneifolia</i> - 1	
<i>Cacoecimorphapronubana</i>	1997 2003	2	<i>Laurus</i> sp. - 1, <i>Ligustrum</i> sp. - 1	
<i>Ailemma loeflingianum</i>	2003	1	<i>Q. castaneifolia</i> - 1	
<i>Tortrix viridana</i>	1994, 1995, 1996, 1997, 1998, 2001, 2002, 2004	12	<i>Quercus</i> sp. - 2, <i>Q. castaneifolia</i> - 251, <i>Tilia</i> sp. - 3, <i>Z. carpinifolia</i> - 30, <i>Mespilus</i> sp. - 11, <i>U. suberosa</i> - 50, <i>P. persica</i> - 40, <i>Cydonia</i> sp. - 9, <i>Crataegus</i> sp. - 10, <i>C. caucasica</i> - 12, <i>P. hyrcana</i> - 6, <i>A. barbata</i> - 3	<i>Q. castaneifolia</i> <i>U. suberosa</i> <i>P. persica</i> <i>Z. carpinifolia</i>
<i>Croesia forskaleana</i>	2001	1	<i>A. hyrcanum</i> - 8	<i>A. hyrcanum</i>
<i>Eudemis profundana</i>	2003, 2004	1	<i>Q. castaneifolia</i> - 5	<i>Q. castaneifolia</i>
<i>Hedya nubiferana</i>	1995 1998 1997	5	<i>Cydonia</i> sp. - 1, <i>Malus</i> sp. - 3, <i>A. vulgaris</i> - 5, <i>P. domestica</i> - 1, <i>Quercus</i> sp. - 1	<i>Malus</i> sp., <i>A. vulgaris</i>
<i>Spilonota ocellana</i>	1995, 2002	2	<i>Malus</i> sp. - 2, <i>Q. castaneifolia</i> - 1	<i>Malus</i> sp.
<i>Epipotia demarniana</i>	2001	1	<i>A. barbata</i> - 2	<i>A. barbata</i>
<i>Gypsonoma</i> sp.	1996	1	<i>Populus</i> sp. - 4	<i>Populus</i> sp.
<i>Notocelia uddmanniana</i>	2001	1	<i>Rubus</i> L. - 2	
<i>Cydia pomonella</i>	1995, 1996, 1997	3	<i>Malus</i> sp. - 4, <i>Q. castaneifolia</i> - 1, <i>A. vulgaris</i> - 1	<i>Malus</i> sp.